

Multiple myeloma melphalan and prednisolone oral (SUPERSEDED)

ID: 58 v.4 Superseded Essential Medicine List

This protocol has been superseded as triplet therapy appears to be more efficacious than single or doublet therapy. It is not commonly used in clinical practice. ID 61 Multiple myeloma MPT (melphalan prednisolone thalidomide) oral is the preferred protocol.

Patients with myeloma should be considered for inclusion into clinical trials. Link to [ALLG website](#) and [ANZCTR website](#).
Link to [Medical Scientific Advisory Group \(MSAG\) Clinical Practice Guideline Multiple Myeloma](#)

The anticancer drug(s) in this protocol may have been included in the ADDIKD guideline. Dose recommendations in kidney dysfunction have yet to be updated to align with the ADDIKD guideline. Recommendations will be updated once the individual protocol has been evaluated by the reference committee. For further information refer to the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\) calculator](#).

International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADDITION)

2022

[Click here](#)



Treatment schedule - Overview

Cycle 1 to 6

Drug	Dose	Route	Day
Prednisolone	60 mg/m ² ONCE a day (Cap dose at 100 mg)	PO	1 to 4
Melphalan	8 mg/m ² ONCE a day	PO	1 to 4

Frequency: 28 days

Cycles: 6 to 8 cycles depending on response. Discontinue when plateau phase achieved or if there is no response by cycle 3. Cycle frequency may be 28 to 35 days.

Notes:

- Consider starting melphalan at a 50% dose reduction for patients >75 years.
- Melphalan has erratic absorption, if no myelosuppression occurs after oral dosing, poor oral absorption should be suspected.
- Several variations exist of the combination melphalan and prednisone regimen.
- Administer with caution within 4 weeks of radiation therapy.

Drug status: Melphalan and prednisolone: [PBS general schedule](#)

Melphalan is available as **2 mg** tablets

Prednisolone is available as **25 mg, 5 mg** and **1 mg** tablets

Cost: ~ \$110 per cycle

Treatment schedule - Detail

The supportive therapies (e.g. antiemetics, premedications, etc.), infusion times, diluents, volumes and routes of administration, if included, are listed as defaults. They may vary between institutions and can be substituted to reflect individual institutional policy.

Antiemetics if included in the treatment schedule are based upon recommendations from national and international guidelines. These are **defaults only** and may be substituted to reflect individual institutional policy. [Select here for recommended doses of alternative antiemetics.](#)

Cycle 1 to 6

Day 1 to 4		
Prednisolone	60 mg/m ² (PO) (Cap dose at 100 mg)	ONCE a day on days 1 to 4. Take in the morning with food.
Melphalan	8 mg/m ² (PO)	ONCE a day on days 1 to 4. Take on an empty stomach at least half an hour before, or 2 hours after food (cap initial dose at 10 mg).

Frequency: 28 days

Cycles: 6 to 8 cycles depending on response. Discontinue when plateau phase achieved or if there is no response by cycle 3. Cycle frequency may be 28 to 35 days.

Indications and patient population

- Patients with multiple myeloma who are not suitable for high dose therapy or transplant

Clinical information

Caution with oral anti-cancer drugs	Select links for information on the safe prescribing, dispensing and administration of orally administered anti-cancer drugs. Read more about the COSA guidelines and oral anti-cancer therapy
Emetogenicity minimal or low	No routine prophylaxis required. If patients experience nausea and/or vomiting, consider using the low emetogenic risk regimen. Read more about preventing anti-cancer therapy induced nausea and vomiting
Bone modifying agents	Use of a bone modifying agent (BMA) should be considered in all patients with symptomatic myeloma requiring treatment. For patients with newly diagnosed symptomatic myeloma, zoledronic acid, pamidronate or denosumab should be considered for monthly administration (adjust for kidney dysfunction where appropriate) for up to 2 years. A longer duration of therapy may be appropriate (MRC M IX trial). ¹ For more information, please see the following protocols: ID 137 Multiple myeloma zoledronic acid ID 147 Multiple myeloma pamidronate ID 3964 Multiple myeloma denosumab - note denosumab is TGA approved but not PBS reimbursed for this indication.
Bisphosphonates and dental review	Caution should be taken with prolonged use of bisphosphonates due to the risk of osteonecrosis of the jaw (ONJ). A dental review prior to treatment is recommended, and all dental issues treated before the initiation of bisphosphonates. Dental review 6 to 12 monthly during treatment is advisable to minimise risk of ONJ. Concurrent daily oral supplements of calcium 500 mg and vitamin D 400 International Units are recommended. Read more about medication-related osteonecrosis of the jaw (MRONJ)
Corticosteroids	Diabetic patients should monitor their blood glucose levels closely. To minimise gastric irritation, advise patient to take immediately after food. Consider the use of a H2 antagonist or proton pump inhibitor if appropriate. Read more about acute short term effects from corticosteroids

Tumour lysis risk	Assess patient for risk of developing tumour lysis syndrome. Read more about prevention and management of tumour lysis syndrome .
Pneumocystis jirovecii pneumonia (PJP) prophylaxis	Read more about prophylaxis of pneumocystis jirovecii (carinii) in cancer patients
Thromboprophylaxis	Thromboprophylaxis should be considered based on an individual benefit/risk assessment and at clinician discretion. Read more about the prophylaxis of venous thromboembolism (VTE) in multiple myeloma
Blood tests	FBC, EUC, LFTs, LDH, calcium, magnesium, phosphate and BSL at baseline, and prior to each treatment and/or as clinically indicated.
Hepatitis B screening and prophylaxis	Routine screening for HBsAg and anti-HBc is recommended prior to initiation of treatment. Prophylaxis should be determined according to individual institutional policy. Read more about hepatitis B screening and prophylaxis in cancer patients requiring cytotoxic and/or immunosuppressive therapy
Vaccinations	Live vaccines are contraindicated in cancer patients receiving immunosuppressive therapy and/or who have poorly controlled malignant disease. Refer to the recommended schedule of vaccination for immunocompromised patients, as outlined in the Australian Immunisation Handbook . Read more about COVID-19 vaccines and cancer .
Fertility, pregnancy and lactation	Cancer treatment can have harmful effects on fertility and this should be discussed with all patients of reproductive potential prior to commencing treatment. There is a risk of foetal harm in pregnant women. A pregnancy test should be considered prior to initiating treatment in females of reproductive potential if sexually active. It is important that all patients of reproductive potential use effective contraception whilst on therapy and after treatment finishes. Effective contraception methods and adequate contraception timeframe should be discussed with all patients of reproductive potential. Possibility of infant risk should be discussed with breastfeeding patients. Read more about the effect of cancer treatment on fertility

Dose modifications

Evidence for dose modifications is limited, and the recommendations made on eviQ are intended as a guide only. They are generally conservative with an emphasis on safety. Any dose modification should be based on clinical judgement, and the individual patient's situation including but not limited to treatment intent (curative vs palliative), the anti-cancer regimen (single versus combination therapy versus chemotherapy versus immunotherapy), biology of the cancer (site, size, mutations, metastases), other treatment related side effects, additional co-morbidities, performance status and patient preferences. Suggested dose modifications are based on clinical trial findings, product information, published guidelines and reference committee consensus. The dose reduction applies to each individual dose and not to the total number of days or duration of treatment cycle unless stated otherwise. Non-haematological gradings are based on [Common Terminology Criteria for Adverse Events \(CTCAE\)](#) unless otherwise specified. Renal and hepatic dose modifications have been standardised where possible. For more information see dosing considerations & disclaimer.

The dose recommendations in kidney dysfunction (i.e. renal impairment) displayed may not reflect those in the ADDIKD guideline and have been included for historical reference only. Recommendations will be updated once the individual protocol has been evaluated by the reference committee, with this version of the protocol then being archived. Clinicians are expected to refer to the ADDIKD guideline prior to prescribing in kidney dysfunction.

[International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction \(ADDIKD\).](#)

Note: All dose reductions are calculated as a percentage of the starting dose

Haematological toxicity

Myelosuppression from melphalan is cumulative from which recovery can be prolonged or incomplete.

Haematological toxicity	
Neutrophils less than $1.0 \times 10^9/L$ and/or platelets less than $75 \times 10^9/L$	Delay melphalan until recovery unless due to marrow infiltration and consider dose reduction on subsequent cycles

Renal impairment	
Creatinine clearance (mL/min)	
30 to 50	Reduce melphalan by 25%
less than 30	Clinical decision but not recommended

Hepatic impairment	
No specific dose modifications recommended for melphalan. If excessive toxicity, consider dose reduction on subsequent cycles.	

Age	
Older than 75 years	Reduce melphalan dose by 50%

Interactions

Drug interactions in eviQ protocols are under review and being updated to align with current literature. Further site-wide updates and changes will occur in due course. *References & Disclaimer*

The drug interactions shown below are not an exhaustive list. For a more comprehensive list and for detailed information on specific drug interactions and clinical management, please refer to the specific drug product information and the following key resources:

- [MIMS - interactions tab](#) (includes link to a CYP-450 table) (login required)
- [Australian Medicines Handbook \(AMH\) – interactions tab](#) (login required)
- [Micromedex Drug Interactions](#) (login required)
- [Cancer Drug Interactions](#)
- [Cytochrome P450 Drug Interactions](#)

Melphalan (oral)		
	Interaction	Clinical management
Ciclosporin	Additive nephrotoxicity	Monitor renal function and ciclosporin levels: dose reduction of ciclosporin may be necessary when used with high dose melphalan

Prednisolone		
	Interaction	Clinical management
Antidiabetic agents (e.g. insulin, glibenclamide, glicazide, metformin, pioglitazone, etc)	The efficacy of antidiabetic agents may be decreased	Use with caution and monitor blood glucose
Azole antifungals (e.g. fluconazole, itraconazole, ketoconazole, posaconazole)	Increased toxicity of prednisolone possible due to reduced clearance	Avoid combination or monitor for prednisolone toxicity
Oestrogens (e.g. oral contraceptives)	Increased toxicity of prednisolone possible due to reduced clearance	Avoid combination or monitor for prednisolone toxicity. Dose reduction of prednisolone may be required
Ritonavir	Increased toxicity of prednisolone possible due to reduced clearance	Avoid combination or monitor for prednisolone toxicity

General		
	Interaction	Clinical management
Warfarin	Anti-cancer drugs may alter the anticoagulant effect of warfarin.	Monitor INR regularly and adjust warfarin dosage as appropriate; consider alternative anticoagulant.
Direct oral anticoagulants (DOACs) e.g. apixaban, rivaroxaban, dabigatran	<p>Interaction with both CYP3A4 and P-gp inhibitors /inducers.</p> <p>DOAC and anti-cancer drug levels may both be altered, possibly leading to loss of efficacy or toxicity (i.e. increased bleeding).</p>	<p>Apixaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. If treating VTE, avoid use with strong CYP3A4 and P-gp inducers.</p> <p>Rivaroxaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors.</p> <p>Dabigatran: avoid combination with strong P-gp inducers and inhibitors.</p> <p>If concurrent use is unavoidable, monitor closely for efficacy/toxicity of both drugs.</p>
Digoxin	Anti-cancer drugs can damage the lining of the intestine; affecting the absorption of digoxin.	Monitor digoxin serum levels; adjust digoxin dosage as appropriate.
Antiepileptics	Both altered antiepileptic and anti-cancer drug levels may occur, possibly leading to loss of efficacy or toxicity.	Where concurrent use of an enzyme-inducing antiepileptic cannot be avoided, monitor antiepileptic serum levels for toxicity, as well as seizure frequency for efficacy; adjust dosage as appropriate. Also monitor closely for efficacy of the anti-cancer therapy.
Antiplatelet agents and NSAIDs	Increased risk of bleeding due to treatment related thrombocytopenia.	Avoid or minimise combination. If combination deemed essential, (e.g. low dose aspirin for ischaemic heart disease) monitor for signs of bleeding.
Serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs e.g. paroxetine) and serotonin noradrenaline reuptake inhibitors (SNRIs e.g. venlafaxine)	Increased risk of serotonin syndrome with concurrent use of 5-HT3 receptor antagonists (e.g. palonosetron, ondansetron, granisetron, tropisetron, dolasetron, etc.)	<p>Avoid combination.</p> <p>If combination is clinically warranted, monitor for signs and symptoms of serotonin syndrome (e.g. confusion, agitation, tachycardia, hyperreflexia).</p> <p>For more information link to TGA Medicines Safety Update</p>
Vaccines	Diminished response to vaccines and increased risk of infection with live vaccines.	<p>Live vaccines (e.g. BCG, MMR, zoster and varicella) are contraindicated in patients on immunosuppressive therapy. Use with caution in patients on non-immunosuppressive therapy.</p> <p>For more information; refer to the recommended schedule of vaccination for cancer patients, as outlined in the Australian Immunisation Handbook</p>

Administration

eviQ provides safe and effective instructions on how to administer cancer treatments. However, eviQ does not provide every treatment delivery option, and is unable to provide a comprehensive list of cancer treatment agents and their required IV line giving set/filter. There may be alternative methods of treatment administration, and alternative supportive treatments that are also appropriate. Please refer to the individual

Days 1 to 4

This is an oral treatment

[Safe handling and waste management](#)

[Safe administration](#)

[General patient assessment](#) prior to each treatment.

Any toxicity grade 2 or greater may require dose reduction, delay or omission of treatment and review by medical officer before recommencing treatment.

Prednisolone

- administer orally ONCE daily
- to be taken in the morning with or immediately after food

Note: missed doses should not be replaced; if a tablet is forgotten or vomited, normal dosing should be resumed at the next scheduled dose.

Chemotherapy - Time out

Melphalan

- administer orally ONCE a day
- to be swallowed whole with a glass of water; do not break, crush or chew
- to be taken on an empty stomach, one hour before or two hours after food
- store tablets in the fridge (2 to 8 degrees C).

Note: if a dose is forgotten or vomited, contact treating team.

Continue [safe handling](#) precautions until 7 days after completion of drug(s)

Discharge information

Melphalan and prednisolone tablets

- Melphalan and prednisolone tablets with written instructions on how to take them.

Antiemetics

- Antiemetics as prescribed.

Prophylaxis medications

- Prophylaxis medications (if prescribed) i.e. tumour lysis prophylaxis, PJP prophylaxis, antifungals, antivirals.

Patient information

- Ensure patient receives patient information sheet.

Side effects

The side effects listed below are not a complete list of all possible side effects for this treatment. Side effects are categorised into the approximate onset of presentation and should only be used as a guide.

Immediate (onset hours to days)

Nausea and vomiting

Read more about [prevention of treatment induced nausea and vomiting](#)

Early (onset days to weeks)	
Neutropenia	Abnormally low levels of neutrophils in the blood. This increases the risk of infection. Any fever or suspicion of infection should be investigated immediately and managed aggressively. Read more about immediate management of neutropenic fever
Thrombocytopenia	A reduction in the normal levels of functional platelets, increasing the risk of abnormal bleeding. Read more about thrombocytopenia
Diarrhoea	Read more about treatment induced diarrhoea
Fatigue	Read more about fatigue
Side effects of corticosteroids	Insomnia, oedema, increased risk of infection e.g. oral thrush, gastric irritation, worsening of peptic ulcer disease, increased blood sugar levels, loss of diabetic control, mood and behavioural changes - including anxiety, euphoria, depression, mood swings, increased appetite and weight gain, osteoporosis and fractures (long term use), bruising and skin fragility are associated with corticosteroid use.
Thromboembolism	Thromboembolic events, including pulmonary embolism, deep vein thrombosis and cerebrovascular accidents can occur. Thromboprophylaxis should be considered based on an individual benefit/risk assessment and at clinician discretion. Read more about management of thromboembolism (VTE) in multiple myeloma

Late (onset weeks to months)	
Anaemia	Abnormally low levels of red blood cells (RBCs) or haemoglobin in the blood. Read more about anaemia
Alopecia - partial	Hair thinning and/or patchy hair loss. Patients can also experience mild to moderate discomfort of the hair follicles, and rarely pain as the hair is falling out. Read more about alopecia and scalp cooling

Evidence

This protocol has been superseded as triplet therapy appears to be more efficacious than single or doublet therapy.

Melphalan and prednisolone (MP) has been the standard induction therapy for myeloma since its introduction in 1958. Approximately 40% of newly diagnosed myeloma patients will achieve a response (defined by at least a 75% reduction in serum paraprotein, 95% reduction in Bence-Jones proteinuria, less than 5% plasma cells on bone marrow biopsy).²

Since 1958, numerous studies comparing MP with various combinations of alkylating agents have been performed. In 1998, the Myeloma Trialists Collaborative Group evaluated 6,633 patients from 27 randomised trials comparing combination chemotherapy with MP. Although the median overall response rate was 60% for the various combination schedules versus 53.2% for MP ($p<0.00001$), there were no significant differences in overall survival. The median survival for both groups was 29 months.³

Efficacy

Results from the Myeloma Trialists' Collaborative Group,³ comparing MP to combination chemotherapy (CCT).

	RR %	Median Survival	5 yr OS
MP	53.2 %	29 months	23%
CCT	60%	29 months	24.4%
p-value	<0.0001	>0.1	>0.1

Response rates were higher with combination chemotherapy treatment (CCT) than with M/P (60.0% v 53.2%; $p<0.0001$, two tailed). There was no evidence of any difference in mortality between CCT and MP with a non-significant 1.5% reduction in death rate in favour of CCT ($p=0.6$, two tailed).

There is heterogeneity of design between the trials, but subgroup analysis by type of CCT or by dose-intensities of CCT, of melphalan, or of prednisone did not identify any particular forms of therapy that were either clearly beneficial or clearly adverse.

This overview found no difference, either overall or within any subgroup, in mortality between CCT and MP. In terms of survival, these therapeutic options, as tested in the trials considered, are approximately equivalent.

Toxicity

Melphalan and prednisone (MP) are generally well tolerated and alopecia is rare. Mild degrees of nausea occur, as do skin rash and amenorrhoea.

Myelosuppression is cumulative and can be prolonged. Nadir 10 to 21 days and recovery 18 to 40 days. Secondary malignancies have been reported (acute non-lymphocytic leukaemia). A summary of the major toxicities is tabulated below.

Toxicity ⁴	Grade	Incidence (%) (n=221)
Neutropenia	≥3	37
Thrombocytopenia	≥3	23
Nausea and vomiting	≥2	10
Infection	≥3	14

© Cancer 1997

References

- 1 Morgan, G. J., J. A. Child, W. M. Gregory, et al. 2011. "Effects of zoledronic acid versus clodronic acid on skeletal morbidity in patients with newly diagnosed multiple myeloma (MRC Myeloma IX): secondary outcomes from a randomised controlled trial." *Lancet Oncol* 12(8):743-752.
- 2 Alexanian, R. and M. Dimopoulos. 1994. "The treatment of multiple myeloma." *N.Engl.J.Med.* 330(7):484-489.
- 3 1998. "Combination chemotherapy versus melphalan plus prednisone as treatment for multiple myeloma: an overview of 6,633 patients from 27 randomized trials. Myeloma Trialists' Collaborative Group." *J.Clin Oncol.* 16(12):3832-3842.
- 4 Oken, M. M., D. P. Harrington, N. Abramson, et al. 1997. "Comparison of melphalan and prednisone with vincristine, carmustine, melphalan, cyclophosphamide, and prednisone in the treatment of multiple myeloma: results of Eastern Cooperative Oncology Group Study E2479." *Cancer.* 79(8):1561-1567.

Bibliography

Durie, B. G., R. A. Kyle, A. Belch, et al. 2003. "Myeloma management guidelines: a consensus report from the Scientific Advisors of the International Myeloma Foundation." *Hematol.J.* 4(6):379-398.

British Committee for standards in Haematology: Guidelines on the diagnosis and management of Multiple Myeloma. UKMF website 2001

Oken MM, Pomeroy C, Weisdorf D, et al. 1996 Prophylactic antibiotics for the prevention of early infection in multiple myeloma. *Am J Med Jun*;100(6):624-8.

History

Version 4

Date	Summary of changes
04/07/2007	Reformatting and addition of extra information.
03/08/2008	Addition of extra information, including the use of routine bisphosphonates; information on Hepatitis B reactivation and PCP.
07/09/2009	Reviewed and transferred to eviQ.
19/01/2012	New format to allow for export of protocol information.

Date	Summary of changes
	Protocol version number changed to v.2. Antiemetics and premedications added to the treatment schedule. Additional Clinical Information, Key Prescribing table and Key Administration table combined into new section titled Clinical Considerations. Drug specific information placed behind the drug name link.
20/01/2012	PHC view added.
31/08/2012	Protocol reviewed using the stratified review process at the Haematology Reference Committee meeting. No change and next review in 2 years.
19/06/2013	Added new bisphosphonate clinical information block.
13/03/2015	Discussed at Haematology Reference Committee meeting. MPT is superior to MT. Supersede protocol.
31/05/2017	Transferred to new eviQ website. Version number change to v.4.
24/05/2019	Reviewed at Haematology Reference Committee meeting. Consensus to remain superseded.
21/12/2021	Changed antiemetic clinical information block to minimal or low, to align with new categories. See ID 7 Prevention of anti-cancer therapy induced nausea and vomiting (AINV) v5.
14/10/2022	The following changes have been made with the consensus agreement of the Haematology Reference Committee: <ul style="list-style-type: none"> • Bone modifying agents block added to "Clinical information" section, related note removed from treatment schedule and linked pages removed • Link to Medical Scientific Advisory Group (MSAG) guidelines updated • Thromboprophylaxis information added to "Clinical information" section • Thromboembolism side effect updated

The information contained in this protocol is based on the highest level of available evidence and consensus of the eviQ reference committee regarding their views of currently accepted approaches to treatment. Any clinician (medical oncologist, haematologist, radiation oncologist, medical physicist, radiation therapist, pharmacist or nurse) seeking to apply or consult this protocol is expected to use independent clinical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use is subject to eviQ's disclaimer available at www.eviQ.org.au

First approved: 31 August 2012
Last reviewed: 24 May 2019
Review due: 24 May 2021
Superseded: 13 March 2015

The currency of this information is guaranteed only up until the date of printing, for any updates please check:

<https://www.eviq.org.au/p/58>

12 Jun 2023

Patient information - Multiple myeloma - Melphalan and prednisolone

Patient's name:

Your treatment

The treatment schedule below explains how the drugs for this treatment are given.


Melphalan and prednisolone		
This treatment cycle is repeated every 28 to 35 days. Your doctor will advise you of the number of treatments you will have.		
Day	Treatment	How it is given
1 to 4	Prednisolone (<i>pred-NIS-oh-lone</i>)	Take orally ONCE a day in the morning with food on days 1 to 4 only.
1 to 4	Melphalan (<i>MEL-fa-lan</i>)	Take orally ONCE a day in the morning on days 1 to 4 only. Take on an empty stomach, at least one hour before or two hours after food. Swallow whole with a glass of water, do not break, chew or crush. Melphalan tablets need to be stored in the fridge.

Missed doses:

- **Melphalan** and **prednisolone**: if you forget to take your tablets or vomit your tablets, contact your treating team.

When to get help

Anticancer drugs (drugs used to treat cancer) can sometimes cause serious problems. It is important to get medical help immediately if you become unwell.

 IMMEDIATELY go to your nearest hospital Emergency Department, or contact your doctor or nurse if you have any of the following at any time:	Emergency contact details Ask your doctor or nurse from your treating team who to contact if you have a problem
<ul style="list-style-type: none">• a temperature of 38°C or higher• chills, sweats, shivers or shakes• shortness of breath• uncontrolled vomiting or diarrhoea• pain, tingling or discomfort in your chest or arms• you become unwell.	Daytime:..... Night/weekend:..... Other instructions:.....

Other information about your treatment

Changes to your dose or treatment delays

Sometimes a treatment may be started at a lower dose or the dose needs to be changed during treatment. There may also be times when your treatment is delayed. This can happen if your doctor thinks you are likely to have severe side effects, if you get severe side effects, if your blood counts are affected and causing delays in treatment, or if you are finding it hard to cope with the treatment. This is called a dose reduction, dose change or treatment delay. Your doctor will explain if you need any changes or delays to your treatment and the reason why.

Blood tests and monitoring

You will need to have a blood test before you start treatment and regularly throughout your treatment. Your doctor or nurse will tell you when to have these blood tests.

Other medications given during this treatment

- **Anti-sickness (anti-nausea) medication:** you may be given some anti-sickness medication. Make sure you take this medication as your doctor or nurse tells you, even if you don't feel sick. This can help to prevent the sickness starting.
- **Prophylaxis medication:** you may need to take some medications to prevent infection and to help prevent or reduce some of the side effects of the chemotherapy. Your doctor or nurse will tell you how and when to take these medications.

Superseded treatments

This treatment is superseded meaning that better treatments have taken its place. Uncommonly superseded treatments are still used. Your doctor will explain why this treatment has been selected for you.

Side effects

Cancer treatments can cause damage to normal cells in your body, which can cause side effects. Everyone gets different side effects, and some people will have more problems than others.

The table below shows some of the side effects you may get with this treatment. You are unlikely to get all of those listed and you may also get some side effects that have not been listed.

Tell your doctor or nurse about any side effects that worry you. Follow the instructions below and those given to you by your doctor or nurse.

Immediate (onset hours to days)	
Nausea and vomiting	<ul style="list-style-type: none">• You may feel sick (nausea) or be sick (vomit).• Take your anti-sickness medication as directed even if you don't feel sick.• Drink plenty of fluids (unless you are fluid restricted).• Eat small meals more frequently.• Try food that does not require much preparation.• Try bland foods like dry biscuits or toast.• Gentle exercise may help with nausea.• Ask your doctor or nurse for eviQ patient information - Nausea and vomiting during cancer treatment.• Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have uncontrolled vomiting or feel dizzy or light-headed.
Early (onset days to weeks)	

Infection risk (neutropenia)	<ul style="list-style-type: none"> • This treatment lowers the amount of white blood cells in your body. The type of white blood cells that help to fight infection are called neutrophils. Having low level of neutrophils is called neutropenia. If you have neutropenia, you are at greater risk of getting an infection. It also means that your body can't fight infections as well as usual. This is a serious side effect, and can be life threatening. • Wash your hands often. • Keep a thermometer at home and take your temperature regularly, and if you feel unwell. • Do your mouth care regularly. • Inspect your central line site (if you have one) daily for any redness, pus or swelling. • Limit contact with people who are sick. • Learn how to recognise the signs of infection. • Ask your doctor or nurse for eviQ patient information - Infection during cancer treatment. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the following signs or symptoms: <ul style="list-style-type: none"> ◦ a temperature of 38°C or higher ◦ chills, shivers, sweats or shakes ◦ a sore throat or cough ◦ uncontrolled diarrhoea ◦ shortness of breath ◦ a fast heartbeat ◦ become unwell even without a temperature.
Low platelets (thrombocytopenia)	<ul style="list-style-type: none"> • This treatment lowers the amount of platelets in your blood. Platelets help your blood to clot. When they are low, you are at an increased risk of bleeding and bruising. • Try not to bruise or cut yourself. • Avoid contact sport or vigorous exercise. • Clear your nose by blowing gently. • Avoid constipation. • Brush your teeth with a soft toothbrush. • Don't take aspirin, ibuprofen or other similar anti-inflammatory medications unless your doctor tells you to. • Tell your doctor or nurse if you have any bruising or bleeding. • Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if you have any uncontrolled bleeding.
Diarrhoea	<ul style="list-style-type: none"> • You may get bowel motions (stools, poo) that are more frequent or more liquid. • You may also get bloating, cramping or pain. • Take your antidiarrhoeal medication as directed by your doctor. • Drink plenty of fluids (unless you are fluid restricted). • Eat and drink small amounts more often. • Avoid spicy foods, dairy products, high fibre foods, and coffee. • Ask your doctor or nurse for eviQ patient information - Diarrhoea during cancer treatment. • Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if your diarrhoea is not controlled, you have 4 or more loose bowel motions per day, and if you feel dizzy or light-headed.
Tiredness and lack of energy (fatigue)	<ul style="list-style-type: none"> • You may feel very tired, have no energy, sleep a lot, and not be able to do normal activities or things you enjoy. • Do not drive or operate machinery if you are feeling tired. • Nap for short periods (only 1 hour at a time) • Prioritise your tasks to ensure the best use of your energy. • Eat a well balanced diet and drink plenty of fluids (unless you are fluid restricted). • Try some gentle exercise daily. • Allow your friends and family to help. • Tell your doctor or nurse if you get any of the symptoms listed above.

Side effects from steroid medication	<ul style="list-style-type: none"> • Steroid medication may cause: <ul style="list-style-type: none"> ◦ mood swings and behaviour changes ◦ an increased appetite ◦ weight gain ◦ swelling in your hands and feet ◦ stomach upsets ◦ trouble sleeping ◦ fragile skin and bruising ◦ an increase in your blood sugar level ◦ weak and brittle bones (osteoporosis) • Take your steroid medication with food to reduce stomach upset • If you have diabetes, your blood sugar levels may be tested more often. • Tell your doctor or nurse if you get any of the symptoms listed above.
---	---

Late (onset weeks to months)	
Low red blood cells (anaemia)	<ul style="list-style-type: none"> • You may feel dizzy, light-headed, tired and appear more pale than usual. • Tell your doctor or nurse if you have any of these signs or symptoms. You might need a blood transfusion. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have any chest pain, trouble breathing, or feel like your heart is racing.
Hair thinning	<ul style="list-style-type: none"> • Your hair may become dry and may break easily. • You may lose some of your hair. • Use a gentle shampoo and a soft hairbrush. • Take care with hair products like hairspray, hair dye, bleaches and perms. • Protect your scalp from the cold with a hat or scarf. • Protect your scalp from the sun with a hat and sunscreen of SPF 50 or higher. • Ask your doctor or nurse about the Look Good Feel Better program (www.lgfb.org.au)

General advice for people having cancer treatment

Chemotherapy safety

- Learn how to keep you and your family safe while you are having anticancer drugs.
- See our patient information sheet - [Chemotherapy safety at home](#).

Blood clot risk

- Cancer and anticancer drugs can increase the risk of a blood clot (thrombosis).
- Tell your doctor if you have a family history of blood clots.
- A blood clot can cause pain, redness, swelling in your arms or legs, shortness of breath or chest pain.
- If you have any of these symptoms go to your nearest hospital Emergency Department.

Medications and vaccinations

- Before you start treatment, tell your doctor about any medications you are taking, including vitamins or herbal supplements.
- Don't stop or start any medications during treatment without talking to your doctor and pharmacist first.
- Paracetamol is safe to take if you have a headache or other mild aches and pains. It is recommended that you avoid taking aspirin, ibuprofen and other anti-inflammatory type medications for pain while you are having treatment. However, if these medications have been prescribed by your doctor, do not stop taking them without speaking with your doctor.
- Vaccinations such as flu and tetanus vaccines are safe to receive while having treatment. Do not have any live vaccines during your treatment or for 6 months after it finishes. If you are unsure, check with your doctor before you have any vaccinations.
- People you live with should be fully vaccinated, including having live vaccines according to the current vaccination schedule. Extra care needs to be taken with hand washing and careful disposal of soiled nappies for infants who have recently received the rotavirus vaccine.

Other medical and dental treatment

- If you go to hospital or any other medical appointment (including dental appointments), always tell the person treating you that

you are receiving anticancer drugs.

- Before you have any dental treatment, talk to your doctor.

Diet and food safety

- While you are receiving this treatment it is important that you try to maintain a healthy diet.
- Speak to your doctor or nurse about whether drinking alcohol is safe with your treatment.
- If you have any concerns about recent weight loss or weight gain or questions about your diet, ask to speak to a dietitian.
- There are some foods that may cause infection in high risk individuals and should be avoided. For more information on foods to avoid and food hygiene please ask for a copy of the [Listeria and food brochure](#).

Fertility

- Some cancer treatments can reduce your fertility. This can make it difficult or impossible to get pregnant or father a child.
- Talk to your doctor or nurse before you start any treatment. Depending on your situation there may be fertility sparing options available to you and/or your partner, discuss these with your doctor or nurse.

Pregnancy and breastfeeding

- Some cancer treatments can be dangerous to unborn babies. Talk to your doctor or nurse if you think there is any chance that you could be pregnant.
- Do not try to get pregnant or father a child during this treatment. Contraception should be used during treatment and after stopping treatment. Ask your doctor or nurse about what type of contraception you should use.
- If you are planning pregnancy/fatherhood after completing this treatment, talk to your doctor. Some doctors advise waiting between 6 months and 2 years after treatment.
- Do not breastfeed if you are on this treatment, as anti-cancer medications can also pass into breast milk.

Sex life and sexuality

- The desire to have sex may decrease as a result of this treatment or its side effects.
- Your emotions and the way you feel about yourself may also be affected by this treatment.
- It may help to discuss your concerns with your partner and doctor or nurse.

Risk of developing a second cancer

- Some anticancer treatments can increase your chance of developing a second cancer, this is rare. Your doctor will discuss with you the specific risks of your treatment.

Quitting smoking

- It is never too late to quit smoking. Quitting smoking is one of the best things you can do to help your treatment work better.
- There are many effective tools to improve your chances of quitting.
- Talk to your treating team for more information and referral to a smoking cessation support service.

Staying active

- Research shows that exercise, no matter how small, has many benefits for people during and after cancer treatment.
- Talk to your doctor before starting an exercise program. Your doctor can advise whether you need a modified exercise program.

For more information about cancer treatment, side effects and side effect management see our [Patient and carers](#) section.

Where to get more information

Telephone support

- Call Cancer Council on 13 11 20 for cancer information and support
- Call the Leukaemia Foundation on 1800 620 420 (Mon to Fri 9am – 5pm)
- Call the Lymphoma Nurse Support Line on 1800 953 081 (Mon to Fri 9am - 5pm)

Haematology, transplant and cellular therapy information

- Arrow bone marrow transplant foundation – arrow.org.au
- Australasian Menopause Society – menopause.org.au
- Chris O'Brien Lifehouse - Total Body Irradiation - mylifehouse.org.au/departments/radiation-oncology/total-body-irradiation/
- Healthy Male Andrology Australia – healthymale.org.au/

- International Myeloma Foundation – myeloma.org
- Leukaemia Foundation – leukaemia.org.au
- Lymphoma Australia – lymphoma.org.au
- Myeloma Australia – myeloma.org.au
- NSW Agency for Clinical Innovation, Blood & Marrow Transplant Network – aci.health.nsw.gov.au/resources/blood-and-marrow-transplant
- NSW Agency for Clinical Innovation - aci.health.nsw.gov.au/projects/immune-effector-cell-service
- NCCN Guidelines for Patients Immunotherapy Side Effects: CAR T-Cell Therapy - nccn.org/patientresources/patient-resources/guidelines-for-patients
- Talk Blood Cancer – cmlsupport.org.uk/organisation-type/social-media-groups

General cancer information and support

- Australian Rare Cancer (ARC) Portal – arcportal.org.au/
- Beyondblue – beyondblue.org.au
- Cancer Australia – canceraustralia.gov.au
- Cancer Council Australia – cancer.org.au
- Cancer Voices Australia – cancervoicesaustralia.org
- CanTeen – canteen.org.au
- Carers Australia – carersaustralia.com.au
- eviQ Cancer Treatments Online – eviQ.org.au
- Food Standards Australia New Zealand: Listeria & Food Safety – foodstandards.gov.au/publications/pages/listeriabrochuretext.aspx
- LGBTQI+ People and Cancer - cancercouncil.com.au/cancer-information/lgbtqi
- Look Good Feel Better – lgfb.org.au
- Patient Information - patients.cancer.nsw.gov.au
- Radiation Oncology Targeting Cancer - targetingcancer.com.au
- Redkite – redkite.org.au
- Return Unwanted Medicines – returnmed.com.au
- Staying active during cancer treatment – patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active

Quit smoking information and support

Quitting smoking is helpful even after you have been diagnosed with cancer. The following resources provide useful information and support to help you quit smoking. Talk to your treating team about any other questions you may have.

- Call Quitline on 13 QUIT (13 78 48)
- iCanQuit – iCanQuit.com.au
- Patient Information - patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking
- Quitnow – quitnow.gov.au

Additional notes:

This document is a guide only and cannot cover every possible situation. The health professionals caring for you should always consider your individual situation when making decisions about your care. Contact your cancer clinic staff or doctor if you have any questions or concerns about your treatment, or you are having problems coping with side effects. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use of this document is subject to eviQ's disclaimer available at www.eviq.org.au

First approved: 31 August 2012

Last reviewed: 24 May 2019

Review due: 24 May 2021

Superseded: 13 March 2015

The currency of this information is guaranteed only up until the date of printing, for any updates please check:

<https://www.eviq.org.au/pi/58>

12 Jun 2023