

Breast metastatic CMF IV (CYCLOPHOSPHamide methotrexate fluorouracil) SUPERSEDED

ID: 45 v.4 **Superseded** Essential Medicine List

This protocol has been superseded because it is not considered best practice for this patient population. For more information see note in *Treatment schedule* below.

⚠ Fluoropyrimidine overdose or overexposure:

Fluoropyrimidine overdose or overexposure may result in severe or life-threatening toxicity. An antidote is available and is highly effective if given within 96 hours. Read more about [fluoropyrimidine overdose or overexposure](#).

Check for clinical trials in this patient group. Link to [Australian Clinical Trials](#) website

The anticancer drug(s) in this protocol may have been included in the ADDIKD guideline. Dose recommendations in kidney dysfunction have yet to be updated to align with the ADDIKD guideline. Recommendations will be updated once the individual protocol has been evaluated by the reference committee. For further information refer to the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\) calculator](#).

International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADIKD)

2022

[Click here](#)



Treatment schedule - Overview

Cycle 1 and further cycles

Drug	Dose	Route	Day
Fluorouracil	600 mg/m ²	IV	1
Methotrexate	40 mg/m ²	IV bolus	1
CYCLOPHOSPHamide	600 mg/m ²	IV infusion	1

Frequency: 21 days

Cycles: Continuous until disease progression or unacceptable toxicity; usually up to 6 cycles

Notes:

"Classical CMF" (oral cyclophosphamide 100 mg/m² D1-14, IV 5-fluorouracil 600 mg/m² and IV methotrexate 40 mg/m² D1 and 8, Q 4/52) has been compared with "modified" intravenous CMF (all drugs given IV D1 Q 3/52) in a randomised controlled trial in postmenopausal women with advanced breast cancer. The response rate was significantly better in favour of the classical CMF arm (48% vs 29%, p=0.003), and overall survival was also superior for classical CMF (17 months vs 12 months, p = 0.016). This trial is the only one directly comparing the two CMF regimens. The results are consistent with classical CMF being a superior regimen, probably explained by the higher dose intensity achieved.¹

Based on the results of this study confirming inferiority of the intravenous CMF regimen, it is recommended that IV CMF be considered a superseded regimen, to be replaced by "classical CMF" as the preferred CMF regimen.

Drug status: All drugs in this protocol are on the [PBS general schedule](#)

Cost: ~ \$180 per cycle

Treatment schedule - Detail

The supportive therapies (e.g. antiemetics, premedications, etc.), infusion times, diluents, volumes and routes of administration, if included, are listed as defaults. They may vary between institutions and can be substituted to reflect individual institutional policy.

Antiemetics if included in the treatment schedule are based upon recommendations from national and international guidelines. These are **defaults only** and may be substituted to reflect individual institutional policy. [Select here for recommended doses of alternative antiemetics.](#)

Cycle 1 and further cycles

Day 1		
Dexamethasone	8 mg (PO)	60 minutes before chemotherapy
Palonosetron	0.25 mg (IV bolus)	30 minutes before chemotherapy
Fluorouracil	600 mg/m ² (IV)	over 3 to 5 minutes
Methotrexate	40 mg/m ² (IV bolus)	over 3 to 5 minutes
CYCLOPHOSPHamide	600 mg/m ² (IV infusion)	in 500 mL sodium chloride 0.9% over 30 to 60 minutes
Day 2 and 3		
Dexamethasone	8 mg (PO)	ONCE a day (or in divided doses) with or after food. Note: dexamethasone doses on day 2 and 3 may not be required and may be reduced or omitted at the clinicians discretion. *

* [Link to ID 7 Prevention of chemotherapy induced nausea and vomiting](#)

Frequency: 21 days

Cycles: Continuous until disease progression or unacceptable toxicity; usually up to 6 cycles

Indications and patient population

- Advanced or metastatic breast cancer
 - For use when an anthracycline based regimen is contraindicated.

Clinical information

Safety alert fluoropyrimidines	Fluoropyrimidines can be administered by different routes and schedules with each method having associated increased risk of certain side effects. Fluoropyrimidine overdose or overexposure is a rare but potentially life threatening side effect of this drug class and can occur by any route of administration. An antidote is available and highly effective if given within 96 hours. Read more about the medication safety alert for infusional fluorouracil and fluoropyrimidine overdose or overexposure
Venous access required	IV cannula (IVC) or central venous access device (CVAD) is required to administer this treatment. Read more about central venous access device line selection

Emetogenicity MODERATE	<p>Suggested default antiemetics have been added to the treatment schedule, and may be substituted to reflect institutional policy.</p> <p>For patients with a prior episode of chemotherapy induced nausea or vomiting, a NK1 receptor antagonist is available on the PBS in combination with a 5HT₃ antagonist and steroid.</p> <p>Ensure that patients also have sufficient antiemetics for breakthrough emesis:</p> <p>Metoclopramide 10 mg three times a day when necessary (maximum of 30 mg/24 hours, up to 5 days) OR</p> <p>Prochlorperazine 10 mg PO every 6 hours when necessary.</p> <p>Read more about preventing anti-cancer therapy induced nausea and vomiting</p>
Cardiac toxicity	<p>Cardiac toxicity is a serious complication that can occur during treatment with fluorouracil. Patients treated with fluorouracil, especially those with a prior history of cardiac disease or other risk factors, should be carefully monitored during therapy.</p> <p>Read more about cardiac toxicity associated with anti-cancer drugs</p>
Pleural effusion or ascites	<p>Methotrexate should be used with caution in patients with pleural effusions or ascites, as methotrexate may accumulate in third space fluid compartments.</p>
Dihydropyrimidine dehydrogenase (DPD) enzyme deficiency	<p>Rare, life-threatening toxicities such as mucositis, neutropenia, neurotoxicity and diarrhoea have been reported following administration of fluoropyrimidines (e.g. fluorouracil and capecitabine). Severe unexplained toxicities require investigation prior to continuing with treatment. Testing for DPD enzyme deficiency is available in Australia but not currently reimbursed.</p> <p>Read more about dihydropyrimidine dehydrogenase (DPD) enzyme deficiency</p>
Blood tests	<p>FBC, EUC and LFTs at baseline and prior to each cycle. INR as clinically indicated.</p>
Hepatitis B screening and prophylaxis	<p>Routine screening for HBsAg and anti-HBc is NOT usually recommended for patients receiving this treatment.</p> <p>Read more about hepatitis B screening and prophylaxis in cancer patients requiring cytotoxic and/or immunosuppressive therapy</p>
Vaccinations	<p>Live vaccines are contraindicated in cancer patients receiving immunosuppressive therapy and/or who have poorly controlled malignant disease.</p> <p>Refer to the recommended schedule of vaccination for immunocompromised patients, as outlined in the Australian Immunisation Handbook.</p> <p>Read more about COVID-19 vaccines and cancer.</p>
Fertility, pregnancy and lactation	<p>Cancer treatment can have harmful effects on fertility and this should be discussed with all patients of reproductive potential prior to commencing treatment. There is a risk of foetal harm in pregnant women. A pregnancy test should be considered prior to initiating treatment in females of reproductive potential if sexually active. It is important that all patients of reproductive potential use effective contraception whilst on therapy and after treatment finishes. Effective contraception methods and adequate contraception timeframe should be discussed with all patients of reproductive potential. Possibility of infant risk should be discussed with breastfeeding patients.</p> <p>Read more about the effect of cancer treatment on fertility</p>

Dose modifications

Evidence for dose modifications is limited, and the recommendations made on eviQ are intended as a guide only. They are generally conservative with an emphasis on safety. Any dose modification should be based on clinical judgement, and the individual patient's situation including but not limited to treatment intent (curative vs palliative), the anti-cancer regimen (single versus combination therapy versus chemotherapy versus immunotherapy), biology of the cancer (site, size, mutations, metastases), other treatment related side effects, additional co-morbidities, performance status and patient preferences. Suggested dose modifications are based on clinical trial findings, product information, published guidelines and reference committee consensus. The dose reduction applies to each individual dose and not to the total number of days or duration of treatment cycle unless stated otherwise. Non-haematological gradings are based on [Common Terminology Criteria for Adverse Events \(CTCAE\)](#) unless otherwise specified. Renal and hepatic dose modifications have been

The dose recommendations in kidney dysfunction (i.e. renal impairment) displayed may not reflect those in the ADDIKD guideline and have been included for historical reference only. Recommendations will be updated once the individual protocol has been evaluated by the reference committee, with this version of the protocol then being archived. Clinicians are expected to refer to the ADDIKD guideline prior to prescribing in kidney dysfunction.

International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADDIKD).

Note: all dose reductions are calculated as a percentage of the starting dose

Haematological toxicity	
ANC x 10⁹/L (pre-treatment blood test)	
1.0 to less than 1.5	Refer to local institutional guidelines; it is the view of the expert clinicians that treatment should continue if patient is clinically well
0.5 to less than 1.0	Delay treatment until recovery
less than 0.5	Delay treatment until recovery and reduce cyclophosphamide, methotrexate and fluorouracil by 25% for subsequent cycles
Febrile neutropenia	Delay treatment until recovery and reduce cyclophosphamide, methotrexate and fluorouracil by 25% for subsequent cycles
Platelets x 10⁹/L (pre-treatment blood test)	
75 to less than 100	The general recommendation is to delay, however if the patient is clinically well it may be appropriate to continue treatment; refer to treating team and/or local institutional guidelines
50 to less than 75	Delay treatment until recovery
less than 50	Delay treatment until recovery and reduce cyclophosphamide, methotrexate and fluorouracil by 25% for subsequent cycles
Renal impairment	
Creatinine clearance (mL/min)	
30 to 50	Reduce fluorouracil and cyclophosphamide by 25% and methotrexate by 50%
less than 30	Reduce fluorouracil and cyclophosphamide by 50% and omit methotrexate or withhold chemotherapy
Hepatic impairment	
Hepatic dysfunction	
Mild	No dose modification necessary
Moderate	Reduce fluorouracil by 25%
Severe	Reduce fluorouracil by 50% and reduce methotrexate by 25%
Mucositis and stomatitis	
Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 st occurrence: No dose reduction 2 nd occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil by 25% 3 rd occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil by 50% 4 th occurrence: Withhold chemotherapy
Grade 3 or Grade 4	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 st occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil by 50% 2 nd occurrence: Withhold chemotherapy

Diarrhoea

Grade 2	<p>Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows:</p> <p>1st occurrence: No dose reduction</p> <p>2nd occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil 25%</p> <p>3rd occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil by 50%</p> <p>4th occurrence: Withhold chemotherapy</p>
Grade 3 or Grade 4	<p>Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows:</p> <p>1st occurrence: Reduce cyclophosphamide, methotrexate and fluorouracil by 50%</p> <p>2nd occurrence: Withhold chemotherapy</p>

Interactions

Drug interactions in eviQ protocols are under review and being updated to align with current literature. Further site-wide updates and changes will occur in due course. [References & Disclaimer](#)

The drug interactions shown below are not an exhaustive list. For a more comprehensive list and for detailed information on specific drug interactions and clinical management, please refer to the specific drug product information and the following key resources:

- [MIMS - interactions tab](#) (includes link to a CYP-450 table) (login required)
- [Australian Medicines Handbook \(AMH\) – interactions tab](#) (login required)
- [Micromedex Drug Interactions](#) (login required)
- [Cancer Drug Interactions](#)
- [Cytochrome P450 Drug Interactions](#)

Cyclophosphamide

	Interaction	Clinical management
CYP3A4 inducers (e.g. carbamazepine, phenytoin, phenobarbitone, rifampicin, St John's wort etc.)	Increased toxicity of cyclophosphamide possible due to increased conversion to active (and inactive) metabolites	Avoid combination or monitor for cyclophosphamide toxicity
CYP3A4 inhibitors (e.g. aprepitant, azole antifungals, clarithromycin, erythromycin, grapefruit juice, ritonavir etc.)	Reduced efficacy of cyclophosphamide possible due to decreased conversion to active (and inactive) metabolites	Avoid combination or monitor for decreased clinical response to cyclophosphamide
Amiodarone	Possible additive pulmonary toxicity with high-dose cyclophosphamide (i.e. doses used prior to stem cell transplant; 60 mg/kg daily or 120 to 270 mg/kg over a few days)	Avoid combination or monitor closely for pulmonary toxicity
Allopurinol, hydrochlorothiazide, indapamide	Delayed effect. Increased risk of bone marrow depression; probably due to reduced clearance of active metabolites of cyclophosphamide	Avoid combination, consider alternative antihypertensive therapy or monitor for myelosuppression
Ciclosporin	Reduced efficacy of ciclosporin due to reduced serum concentration	Monitor ciclosporin levels; adjust dosage as appropriate; monitor response to ciclosporin
Suxamethonium	Prolonged apnoea due to marked and persistent inhibition of cholinesterase by cyclophosphamide	Alert the anaesthetist if a patient has been treated with cyclophosphamide within ten days of planned general anaesthesia

Fluorouracil		
	Interaction	Clinical management
Folic acid	Increased toxicity of fluorouracil due to stabilisation of its bond to thymidylate synthetase (folic acid is a precursor of folinic acid/leucovorin)	Advise patients not to take folic acid supplements (inc. multivitamins) around the time of receiving treatment with fluorouracil
Metronidazole, tinidazole	Increased toxicity of fluorouracil due to reduced clearance	Avoid combination or monitor for fluorouracil toxicity
Warfarin and other drugs metabolised by CYP2C9 (e.g. warfarin, phenytoin etc.)	Increased effect/toxicity of these drugs due to inhibition of CYP2C9 by fluorouracil resulting in reduced clearance	Avoid combination or monitor for increased effect/toxicity of these drugs (e.g. for bleeding/elevated INR with warfarin, elevated phenytoin serum levels or signs of toxicity such as ataxia, tremor etc.)
Allopurinol	Reduced efficacy of fluorouracil possible due to reduced conversion to the active metabolites	Avoid combination or monitor for reduced fluorouracil efficacy

Methotrexate		
	Interaction	Clinical management
Ciprofloxacin NSAIDs Probenecid Proton pump inhibitors (e.g. esomeprazole, omeprazole, pantoprazole)	Increased toxicity of methotrexate possible due to reduced clearance	Avoid combination or monitor for methotrexate toxicity Important note: with high-dose methotrexate therapy, many of these drug combinations are <i>contraindicated</i>
Sulphonamides and penicillins (e.g. sulfamethoxazole (in Bactrim[®], Septrin[®]), piperacillin (in Tazocin[®]) etc.)	Increased toxicity of methotrexate possible due to displacement from serum protein binding	Avoid combination or monitor for methotrexate toxicity
Trimethoprim	Increased toxicity of methotrexate possible due to additive antifolate activity	Avoid combination or monitor for methotrexate toxicity
Mercaptopurine	Increased toxicity of mercaptopurine possible due to reduced clearance	Avoid combination or monitor for mercaptopurine toxicity
Nephrotoxic drugs (e.g. aminoglycosides, amphotericin, contrast dye, frusemide, NSAIDs)	Additive nephrotoxicity	Avoid combination or monitor kidney function closely
Hepatotoxic drugs (e.g. azathioprine, leflunomide, retinoids, sulfasalazine)	Additive hepatotoxicity	Avoid combination or monitor liver function closely
Folic acid (e.g. as in multivitamins) Asparaginase (administered immediately prior or concurrently)	Reduced efficacy of methotrexate possible due antagonism of its action	Avoid combination or monitor for decreased clinical response to methotrexate Note: asparaginase administered shortly after methotrexate can enhance its efficacy and reduce its toxicity

General		
	Interaction	Clinical management
Warfarin	Anti-cancer drugs may alter the anticoagulant effect of warfarin.	Monitor INR regularly and adjust warfarin dosage as appropriate; consider alternative anticoagulant.
Direct oral anticoagulants (DOACs) e.g. apixaban, rivaroxaban, dabigatran	Interaction with both CYP3A4 and P-gp inhibitors /inducers. DOAC and anti-cancer drug levels may both be altered, possibly leading to loss of efficacy or toxicity (i.e. increased bleeding).	Apixaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. If treating VTE, avoid use with strong CYP3A4 and P-gp inducers. Rivaroxaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. Dabigatran: avoid combination with strong P-gp inducers and inhibitors. If concurrent use is unavoidable, monitor closely for efficacy/toxicity of both drugs.
Digoxin	Anti-cancer drugs can damage the lining of the intestine; affecting the absorption of digoxin.	Monitor digoxin serum levels; adjust digoxin dosage as appropriate.
Antiepileptics	Both altered antiepileptic and anti-cancer drug levels may occur, possibly leading to loss of efficacy or toxicity.	Where concurrent use of an enzyme-inducing antiepileptic cannot be avoided, monitor antiepileptic serum levels for toxicity, as well as seizure frequency for efficacy; adjust dosage as appropriate. Also monitor closely for efficacy of the anti-cancer therapy.
Antiplatelet agents and NSAIDs	Increased risk of bleeding due to treatment related thrombocytopenia.	Avoid or minimise combination. If combination deemed essential, (e.g. low dose aspirin for ischaemic heart disease) monitor for signs of bleeding.
Serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs e.g. paroxetine) and serotonin noradrenaline reuptake inhibitors (SNRIs e.g. venlafaxine)	Increased risk of serotonin syndrome with concurrent use of 5-HT3 receptor antagonists (e.g. palonosetron, ondansetron, granisetron, tropisetron, dolasetron, etc.)	Avoid combination. If combination is clinically warranted, monitor for signs and symptoms of serotonin syndrome (e.g. confusion, agitation, tachycardia, hyperreflexia). For more information link to TGA Medicines Safety Update
Vaccines	Diminished response to vaccines and increased risk of infection with live vaccines.	Live vaccines (e.g. BCG, MMR, zoster and varicella) are contraindicated in patients on immunosuppressive therapy. Use with caution in patients on non-immunosuppressive therapy. For more information; refer to the recommended schedule of vaccination for cancer patients, as outlined in the Australian Immunisation Handbook

Administration

eviQ provides safe and effective instructions on how to administer cancer treatments. However, eviQ does not provide every treatment delivery option, and is unable to provide a comprehensive list of cancer treatment agents and their required IV line giving set/filter. There may be alternative methods of treatment administration, and alternative supportive treatments that are also appropriate. Please refer to the individual

Day 1

Approximate treatment time: 2 hours

[Safe handling and waste management](#)

[Safe administration](#)

[General patient assessment](#) prior to each day of treatment.

Any toxicity grade 2 or greater may require dose reduction, delay or omission of treatment and review by medical officer before commencing treatment.

Prime IV line(s).

Insert IV cannula or access [TIVAD](#) or [CVAD](#).

Pre treatment medication

Verify dexamethasone taken or administer as prescribed.

Verify antiemetics taken or administer as prescribed.

Chemotherapy - Time out

Fluorouracil

- If using cryotherapy commence ~ 5 minutes prior to administering fluorouracil and continue for ~ 30 minutes post.

Administer fluorouracil (irritant):

- over 3 to 5 minutes
 - via a minibag **OR**
 - by IV bolus via a side port of a freely flowing IV infusion
- flush with ~ 100 mL of sodium chloride 0.9%.

Methotrexate

- over 3 to 5 minutes
 - by IV bolus via a side port of a freely flowing IV infusion **OR**
- over 5 to 15 minutes
 - via a minibag
- flush with ~50 mL of sodium chloride 0.9%

Cyclophosphamide

Administer cyclophosphamide:

- via IV infusion over 30 to 60 minutes
- flush with ~ 50 mL of sodium chloride 0.9%
- rapid infusion can cause dizziness, rhinitis, nausea and perioral numbness. If symptoms develop, slow infusion rate.

Remove IV cannula and/or deaccess [TIVAD](#) or [CVAD](#).

Continue [safe handling](#) precautions until 7 days after completion of drug(s)

Discharge information

Antiemetics

- Antiemetics as prescribed.

Patient information

- Ensure patient receives patient information sheet.

Side effects

The side effects listed below are not a complete list of all possible side effects for this treatment. Side effects are categorised into the approximate onset of presentation and should only be used as a guide.

Immediate (onset hours to days)

Nausea and vomiting	Read more about prevention of treatment induced nausea and vomiting
Taste and smell alteration	Read more about taste and smell changes

Early (onset days to weeks)

Neutropenia	Abnormally low levels of neutrophils in the blood. This increases the risk of infection. Any fever or suspicion of infection should be investigated immediately and managed aggressively. Read more about immediate management of neutropenic fever
Diarrhoea	Read more about treatment induced diarrhoea
Oral mucositis	Erythematous and ulcerative lesions of the gastrointestinal tract (GIT). It commonly develops following chemotherapy, radiation therapy to the head, neck or oesophagus, and high dose chemotherapy followed by a blood and marrow transplant (BMT). Read more about oral mucositis
Photosensitivity	Increased sensitivity to ultraviolet (UV) light resulting in an exaggerated sunburn-like reaction accompanied by stinging sensations and urticaria.
Actinic keratoses flare	Pre-existing actinic keratoses (AKs) can become more inflamed and scaly as a result of immunosuppression. Read more about actinic keratoses flare
Ocular changes	Symptoms may include eye pain, blurred vision, blepharitis, uveitis, optic neuritis, tear duct stenosis, conjunctivitis, hyperlacrimation, watery or dry eyes and photophobia.
Fatigue	Read more about fatigue

Late (onset weeks to months)


Pulmonary toxicity	Pulmonary toxicity may include damage to the lungs, airways, pleura and pulmonary circulation. Read more about pulmonary toxicity associated with anti-cancer drugs
Anaemia	Abnormally low levels of red blood cells (RBCs) or haemoglobin in the blood. Read more about anaemia
Alopecia - partial	Hair thinning and/or patchy hair loss. Patients can also experience mild to moderate discomfort of the hair follicles, and rarely pain as the hair is falling out. Read more about alopecia and scalp cooling

Evidence

CMF, in its range of schedules, is often considered a standard regimen for first line treatment of metastatic breast cancer with response rates between 30 - 60%. A randomised trial was carried out to compare Methotrexate and Mitozantrone (MM) with CMF incorporating QOL and side effects on treatment. Treatment schedule was as follows: CMF = Cyclophosphamide 600 mg/m², Methotrexate 40 mg/m² and Fluorouracil 600 mg/m² every 4/52 for 6 cycles; MM = Methotrexate 30mg/m² and Mitozantrone 6.5 mg/m² every 3/52 for 8 cycles. Objective responses were achieved in 29% with CMF and 15% with MM. *P*=0.07. No statistically significant differences were seen in overall survival (OS) or time to progression (TTP) between the two regimens although a tendency towards a shorter progression time on the MM regimen must be acknowledged.²


Efficacy

Kaplan-Meier curve of overall survival²



© British Journal of Cancer 1999

Kaplan-Meier curve of progression-free survival²



© British Journal of Cancer 1999

Toxicity

Toxicity of CMF vs MM regimens as per WHO grading.²

Toxicity	CMF All grades (%)	MM All grades (%)	CMF Grade 3-4 (%)	MM Grade 3-4 (%)	p-value
Nausea & vomiting	61	44	19	4	0.08
Diarrhoea	18	6	0	0	0.07
Alopecia	31	8	8	0	0.002
Rash	2	2	0	0	0.7
Anaemia	50	32	4	2	0.2

Toxicity	CMF All grades (%)	MM All grades (%)	CMF Grade 3-4 (%)	MM Grade 3-4 (%)	p-value
Leucopenia	70	54	24	0	0.001

References

- Engelsman, E., J. C. Klijn, R. D. Rubens, et al. 1991. "'Classical' CMF versus a 3-weekly intravenous CMF schedule in postmenopausal patients with advanced breast cancer. An EORTC Breast Cancer Co-operative Group Phase III Trial (10808)." *Eur J Cancer* 27(8):966-970.
- Harper-Wynne, C., J. English, L. Meyer, et al. 1999. "Randomized trial to compare the efficacy and toxicity of cyclophosphamide, methotrexate and 5-fluorouracil (CMF) with methotrexate mitoxantrone (MM) in advanced carcinoma of the breast." *Br.J.Cancer*. 81(2):316-322

History

Version 4

Date	Summary of changes
22/01/2007	Superseded after discussion at reference group meeting.
01/08/2009	Transferred to eviQ.
21/06/2012	Full view of protocol created (abbreviated version currently published) and review date set for 2 years.
09/05/2014	Protocol reviewed by email survey. No change and next review in 2 years.
08/04/2016	Protocol reviewed at Medical Oncology Reference Committee meeting. No changes and next review in 2 years.
16/10/2016	Patient information sheet updated to include more fluorouracil toxicity symptom warnings.
31/05/2017	Transferred to new eviQ website. Version number change to V.3. Hepatitis B screening changed to NOT recommended.
03/11/2017	Reviewed by Medical Oncology Reference Committee. No changes. Review in 2 years.
10/05/2018	Haematological dose modifications updated as per consensus of the expert clinician group. Fluoropyrimidine overdose or overexposure warning added. Fluoropyrimidine safety alert added in clinical information. DPD enzyme deficiency wording in clinical information updated. Version number changed to V.4.
22/06/2018	Antiemetics updated to be in line with international guidelines. Note to dexamethasone added.
23/09/2019	Protocol reviewed at Medical Oncology Reference Committee meeting on 30/08/2019. No changes. Next review in 2 years.
13/08/2021	Protocol reviewed at Medical Oncology Reference Committee meeting on 13/08/2021. No changes. Next review in 2 years.

The information contained in this protocol is based on the highest level of available evidence and consensus of the eviQ reference committee regarding their views of currently accepted approaches to treatment. Any clinician (medical oncologist, haematologist, radiation oncologist, medical physicist, radiation therapist, pharmacist or nurse) seeking to apply or consult this protocol is expected to use independent clinical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use is subject to eviQ's disclaimer available at www.eviq.org.au

First approved: 24 February 2005
Last reviewed: 13 August 2021
Review due: 31 December 2023
Superseded: 22 January 2007

The currency of this information is guaranteed only up until the date of printing, for any updates please check:

<https://www.eviq.org.au/p/45>

05 Jul 2023

Patient information - Breast cancer metastatic - CMF IV (cyclophosphamide, methotrexate, fluorouracil)

Patient's name:

Your treatment

The treatment schedule below explains how the drugs for this treatment are given.


CMF (cyclophosphamide, methotrexate, fluorouracil)

This treatment cycle is repeated every 21 days. Your doctor will advise you of the number of treatments you will have

Day	Treatment	How it is given	How long it takes
1	Fluorouracil (<i>Flure-oh-YOOR-a-sill</i>) Methotrexate (<i>meth-o-trex-ate</i>) Cyclophosphamide (<i>SYE-kloe-FOS-fa-mide</i>)	By a drip into a vein	About 2 hours

When to get help

Anticancer drugs (drugs used to treat cancer) can sometimes cause serious problems. It is important to get medical help immediately if you become unwell.

 IMMEDIATELY go to your nearest hospital Emergency Department, or contact your doctor or nurse if you have any of the following at any time:	Emergency contact details Ask your doctor or nurse from your treating team who to contact if you have a problem
<ul style="list-style-type: none">• a temperature of 38°C or higher• chills, sweats, shivers or shakes• shortness of breath• uncontrolled vomiting or diarrhoea• pain, tingling or discomfort in your chest or arms• you become unwell.	Daytime:..... Night/weekend:..... Other instructions:.....

During your treatment immediately tell the doctor or nurse looking after you if you get any of the following problems:

- leaking from the area where the drugs are being given
- pain, stinging, swelling or redness in the area where the drugs are being given or at any injection sites
- a skin rash, itching, feeling short of breath, wheezing, fever, shivers, or feeling dizzy or unwell in any way (allergic reaction).

Other information about your treatment

Changes to your dose or treatment delays

Sometimes a treatment may be started at a lower dose or the dose needs to be changed during treatment. There may also be

times when your treatment is delayed. This can happen if your doctor thinks you are likely to have severe side effects, if you get severe side effects, if your blood counts are affected and causing delays in treatment, or if you are finding it hard to cope with the treatment. This is called a dose reduction, dose change or treatment delay. Your doctor will explain if you need any changes or delays to your treatment and the reason why.

Blood tests and monitoring

Anti-cancer drugs can reduce the number of blood cells in your body. You will need to have regular blood tests to check that your blood cell count has returned to normal. If your blood count is low your treatment may be delayed until it has returned to normal. Your doctor or nurse will tell you when to have these blood tests. Tell your doctor if you are on an anticoagulant (medication used to treat or prevent blood clots) e.g. warfarin. You may need to have additional blood tests.

Treatment with cyclophosphamide

You should drink at least 8 to 10 glasses of fluid (unless you are fluid restricted) for 2 days after treatment with cyclophosphamide. You should also empty your bladder often.

Other medications given during this treatment

- **Anti-sickness (anti-nausea) medication:** you may be given some anti-sickness medication. Make sure you take this medication as your doctor or nurse tells you, even if you don't feel sick. This can help to prevent the sickness starting.

Superseded treatments

This treatment is superseded meaning that better treatments have taken its place. Uncommonly superseded treatments are still used. Your doctor will explain why this treatment has been selected for you.

Side effects

Cancer treatments can cause damage to normal cells in your body, which can cause side effects. Everyone gets different side effects, and some people will have more problems than others.

The table below shows some of the side effects you may get with this treatment. You are unlikely to get all of those listed and you may also get some side effects that have not been listed.

Tell your doctor or nurse about any side effects that worry you. Follow the instructions below and those given to you by your doctor or nurse.

Immediate (onset hours to days)	
Nausea and vomiting	<ul style="list-style-type: none">• You may feel sick (nausea) or be sick (vomit).• Take your anti-sickness medication as directed even if you don't feel sick.• Drink plenty of fluids (unless you are fluid restricted).• Eat small meals more frequently.• Try food that does not require much preparation.• Try bland foods like dry biscuits or toast.• Gentle exercise may help with nausea.• Ask your doctor or nurse for eviQ patient information - Nausea and vomiting during cancer treatment.• Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have uncontrolled vomiting or feel dizzy or light-headed.
Taste and smell changes	<ul style="list-style-type: none">• You may find that food loses its taste or tastes different.• These changes are likely to go away with time.• Do your mouth care regularly.• Chew on sugar-free gum or eat sugar-free mints.• Add flavour to your food with sauces and herbs.• Ask your doctor or nurse for eviQ patient information - Taste and smell changes during cancer treatment.

Early (onset days to weeks)

Infection risk (neutropenia)	<ul style="list-style-type: none"> • This treatment lowers the amount of white blood cells in your body. The type of white blood cells that help to fight infection are called neutrophils. Having low level of neutrophils is called neutropenia. If you have neutropenia, you are at greater risk of getting an infection. It also means that your body can't fight infections as well as usual. This is a serious side effect, and can be life threatening. • Wash your hands often. • Keep a thermometer at home and take your temperature regularly, and if you feel unwell. • Do your mouth care regularly. • Inspect your central line site (if you have one) daily for any redness, pus or swelling. • Limit contact with people who are sick. • Learn how to recognise the signs of infection. • Ask your doctor or nurse for eviQ patient information - Infection during cancer treatment. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the following signs or symptoms: <ul style="list-style-type: none"> ◦ a temperature of 38°C or higher ◦ chills, shivers, sweats or shakes ◦ a sore throat or cough ◦ uncontrolled diarrhoea ◦ shortness of breath ◦ a fast heartbeat ◦ become unwell even without a temperature.
Diarrhoea	<ul style="list-style-type: none"> • You may get bowel motions (stools, poo) that are more frequent or more liquid. • You may also get bloating, cramping or pain. • Take your antidiarrhoeal medication as directed by your doctor. • Drink plenty of fluids (unless you are fluid restricted). • Eat and drink small amounts more often. • Avoid spicy foods, dairy products, high fibre foods, and coffee. • Ask your doctor or nurse for eviQ patient information - Diarrhoea during cancer treatment. • Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if your diarrhoea is not controlled, you have 4 or more loose bowel motions per day, and if you feel dizzy or light-headed.
Mouth pain and soreness (mucositis)	<ul style="list-style-type: none"> • You may have: <ul style="list-style-type: none"> ◦ bleeding gums ◦ mouth ulcers ◦ a white coating on your tongue ◦ pain in the mouth or throat ◦ difficulty eating or swallowing. • Avoid spicy, acidic or crunchy foods and very hot or cold food and drinks. • Try bland and soft foods. • Brush your teeth gently with a soft toothbrush after each meal and at bedtime. If you normally floss continue to do so. • Rinse your mouth after you eat and brush your teeth, using either: <ul style="list-style-type: none"> ◦ 1/4 teaspoon of salt in 1 cup of warm water, or ◦ 1/4 teaspoon of bicarbonate of soda in 1 cup of warm water • Ask your doctor or nurse for eviQ patient information - Mouth problems during cancer treatment. • Tell your doctor or nurse if you get any of the symptoms listed above.
Skin that is more sensitive to the sun (photosensitivity)	<ul style="list-style-type: none"> • After being out in the sun you may develop a rash like a bad sunburn. • Your skin may become red, swollen and blistered. • Avoid direct sunlight. • Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and a sunscreen of SPF 50 or higher. • Tell your doctor or nurse if you get any of the symptoms listed above.

Skin changes	<ul style="list-style-type: none"> • Your skin may become dry, and you may notice changes to areas of your skin that have been exposed to the sun. • Keep your skin moisturised with a cream such as sorbolene or aqueous cream. • Avoid direct sunlight. • Protect your skin from the sun by wearing a wide-brimmed hat, sun-protective clothing, sunglasses and sunscreen of SPF 50 or higher. • Tell your doctor or nurse if you notice any skin changes.
Eye problems	<ul style="list-style-type: none"> • You may get: <ul style="list-style-type: none"> ◦ eye pain ◦ red, sore or swollen eyes ◦ blurred vision ◦ watery or gritty eyes ◦ changes in your eyesight ◦ sensitivity to sunlight. • Protect your eyes from the weather (sun and wind) by wearing sunglasses, especially if you have lost your eyelashes. • Tell your doctor or nurse if you get any of the symptoms listed above. Eye drops may help with your symptoms.
Tiredness and lack of energy (fatigue)	<ul style="list-style-type: none"> • You may feel very tired, have no energy, sleep a lot, and not be able to do normal activities or things you enjoy. • Do not drive or operate machinery if you are feeling tired. • Nap for short periods (only 1 hour at a time) • Prioritise your tasks to ensure the best use of your energy. • Eat a well balanced diet and drink plenty of fluids (unless you are fluid restricted). • Try some gentle exercise daily. • Allow your friends and family to help. • Tell your doctor or nurse if you get any of the symptoms listed above.

Late (onset weeks to months)	
Lung problems	<ul style="list-style-type: none"> • Lung problems are rare, but can be serious. They may occur throughout treatment or after the completion of treatment. • You may get: <ul style="list-style-type: none"> ◦ shortness of breath ◦ fever ◦ dry cough ◦ wheezing ◦ fast heartbeat ◦ chest pain. • Your doctor will monitor how well your lungs are working during your treatment. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have chest pain or become short of breath.
Low red blood cells (anaemia)	<ul style="list-style-type: none"> • You may feel dizzy, light-headed, tired and appear more pale than usual. • Tell your doctor or nurse if you have any of these signs or symptoms. You might need a blood transfusion. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have any chest pain, trouble breathing, or feel like your heart is racing.
Hair thinning	<ul style="list-style-type: none"> • Your hair may become dry and may break easily. • You may lose some of your hair. • Use a gentle shampoo and a soft hairbrush. • Take care with hair products like hairspray, hair dye, bleaches and perms. • Protect your scalp from the cold with a hat or scarf. • Protect your scalp from the sun with a hat and sunscreen of SPF 50 or higher. • Ask your doctor or nurse about the Look Good Feel Better program (www.lgfb.org.au)

Chemotherapy safety

- Learn how to keep you and your family safe while you are having anticancer drugs.
- See our patient information sheet - [Chemotherapy safety at home](#).

Blood clot risk

- Cancer and anticancer drugs can increase the risk of a blood clot (thrombosis).
- Tell your doctor if you have a family history of blood clots.
- A blood clot can cause pain, redness, swelling in your arms or legs, shortness of breath or chest pain.
- If you have any of these symptoms go to your nearest hospital Emergency Department.

Medications and vaccinations

- Before you start treatment, tell your doctor about any medications you are taking, including vitamins or herbal supplements.
- Don't stop or start any medications during treatment without talking to your doctor and pharmacist first.
- Paracetamol is safe to take if you have a headache or other mild aches and pains. It is recommended that you avoid taking aspirin, ibuprofen and other anti-inflammatory type medications for pain while you are having treatment. However, if these medications have been prescribed by your doctor, do not stop taking them without speaking with your doctor.
- Vaccinations such as flu and tetanus vaccines are safe to receive while having treatment. Do not have any live vaccines during your treatment or for 6 months after it finishes. If you are unsure, check with your doctor before you have any vaccinations.
- People you live with should be fully vaccinated, including having live vaccines according to the current vaccination schedule. Extra care needs to be taken with hand washing and careful disposal of soiled nappies for infants who have recently received the rotavirus vaccine.

Other medical and dental treatment

- If you go to hospital or any other medical appointment (including dental appointments), always tell the person treating you that you are receiving anticancer drugs.
- Before you have any dental treatment, talk to your doctor.

Diet

- While you are receiving this treatment it is important that you try to maintain a healthy diet.
- Grapefruit and grapefruit juice can interact with your medication and should be avoided while you are on this treatment.
- Speak to your doctor or nurse about whether drinking alcohol is safe with your treatment.
- If you have any concerns about recent weight loss or weight gain or questions about your diet, ask to speak to a dietitian.

Fertility

- Some cancer treatments can reduce your fertility. This can make it difficult or impossible to get pregnant or father a child.
- Talk to your doctor or nurse before you start any treatment. Depending on your situation there may be fertility sparing options available to you and/or your partner, discuss these with your doctor or nurse.

Pregnancy and breastfeeding

- Some cancer treatments can be dangerous to unborn babies. Talk to your doctor or nurse if you think there is any chance that you could be pregnant.
- Do not try to get pregnant or father a child during this treatment. Contraception should be used during treatment and after stopping treatment. Ask your doctor or nurse about what type of contraception you should use.
- If you are planning pregnancy/fatherhood after completing this treatment, talk to your doctor. Some doctors advise waiting between 6 months and 2 years after treatment.
- Do not breastfeed if you are on this treatment, as anti-cancer medications can also pass into breast milk.

Sex life and sexuality

- The desire to have sex may decrease as a result of this treatment or its side effects.
- Your emotions and the way you feel about yourself may also be affected by this treatment.
- It may help to discuss your concerns with your partner and doctor or nurse.

Risk of developing a second cancer

- Some anticancer treatments can increase your chance of developing a second cancer, this is rare. Your doctor will discuss with you the specific risks of your treatment.

Quitting smoking

- It is never too late to quit smoking. Quitting smoking is one of the best things you can do to help your treatment work better.
- There are many effective tools to improve your chances of quitting.
- Talk to your treating team for more information and referral to a smoking cessation support service.

Staying active

- Research shows that exercise, no matter how small, has many benefits for people during and after cancer treatment.
- Talk to your doctor before starting an exercise program. Your doctor can advise whether you need a modified exercise program.

For more information about cancer treatment, side effects and side effect management see our [Patient and carers](#) section.

Where to get more information

Telephone support

- Call Cancer Council on 13 11 20 for cancer information and support.

Breast cancer information

- Australasian Lymphology Association – lymphoedema.org.au
- Australasian Menopause Society – menopause.org.au
- Breast Cancer Network Australia – bcna.org.au
- National Breast Cancer Foundation – nbcf.org.au
- YWCA Encore breast cancer exercise program – ywcaencore.org.au

General cancer information and support

- Australian Rare Cancer (ARC) Portal – arcportal.org.au/
- Beyondblue – beyondblue.org.au
- Cancer Australia – canceraustralia.gov.au
- Cancer Council Australia – cancer.org.au
- Cancer Voices Australia – cancervoicesaustralia.org
- CanTeen – canteen.org.au
- Carers Australia – carersaustralia.com.au
- CHILL Cancer related hair loss – scalpcooling.org
- eviQ Cancer Treatments Online – eviQ.org.au
- LGBTQI+ People and Cancer - cancercouncil.com.au/cancer-information/lgbtqi
- Look Good Feel Better – lgfb.org.au
- Patient Information – patients.cancer.nsw.gov.au
- Radiation Oncology Targeting Cancer – targetingcancer.com.au
- Redkite – redkite.org.au
- Return Unwanted Medicines – returnmed.com.au
- Staying active during cancer treatment – patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active

Quit smoking information and support

Quitting smoking is helpful even after you have been diagnosed with cancer. The following resources provide useful information and support to help you quit smoking. Talk to your treating team about any other questions you may have.

- Call Quitline on 13 QUIT (13 78 48)
- iCanQuit – iCanQuit.com.au
- Patient Information – patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking
- Quitnow – quitnow.gov.au

Additional notes:

This document is a guide only and cannot cover every possible situation. The health professionals caring for you should always consider your individual situation when making decisions about your care. Contact your cancer clinic staff or doctor if you have any questions or concerns about your treatment, or you are having problems coping with side effects. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use of this document is subject to eviQ's disclaimer available at www.eviq.org.au

First approved: 24 February 2005
Last reviewed: 13 August 2021
Review due: 31 December 2023
Superseded: 22 January 2007

The currency of this information is guaranteed only up until the date of printing, for any updates please check:

<https://www.eviq.org.au/pi/45>

05 Jul 2023