

# Colorectal metastatic FOLFIRI (modified) (fluorouracil leucovorin irinotecan) and cetuximab (two weekly)

ID: 1682 v.5 Endorsed

## ⚠ Fluoropyrimidine overdose or overexposure:

Fluoropyrimidine overdose or overexposure may result in severe or life-threatening toxicity. An antidote is available and is highly effective if given within 96 hours. Read more about [fluoropyrimidine overdose or overexposure](#).

Check for clinical trials in this patient group. Link to [Australian Clinical Trials](#) website

This protocol is based on limited evidence; refer to the evidence section of this protocol for more information.

The anticancer drug(s) in this protocol may have been included in the ADDIKD guideline. Dose recommendations in kidney dysfunction have yet to be updated to align with the ADDIKD guideline. Recommendations will be updated once the individual protocol has been evaluated by the reference committee. For further information refer to the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\) calculator](#).

## International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADDIKD)

2022

[Click here](#)



### Related pages:

- [Colorectal metastatic FOLFIRI \(modified\) \(fluorouracil leucovorin irinotecan\) and cetuximab SUPERSEDED](#)

## Treatment schedule - Overview

### Cycle 1 and further cycles

Drug	Dose	Route	Day
Cetuximab	500 mg/m <sup>2</sup>	IV infusion	1
Irinotecan	180 mg/m <sup>2</sup>	IV infusion	1
Calcium folinate (Leucovorin)	50 mg *	IV bolus	1
Fluorouracil	400 mg/m <sup>2</sup>	IV	1
Fluorouracil	2,400 mg/m <sup>2</sup>	CIV via pump over 46 hours	1

\* The dose of calcium folinate (Leucovorin<sup>®</sup>) has been modified in this protocol from the original clinical trial dose of 200 mg/m<sup>2</sup> to 50 mg based on reference committee consensus. Refer to discussion on calcium folinate (Leucovorin<sup>®</sup>) and evidence section for more information.

**Frequency:** 14 days

**Cycles:** Continuous until disease progression or unacceptable toxicity

**Drug status:** Fluorouracil, leucovorin and irinotecan are on the [PBS general schedule](#) and cetuximab is [PBS authority](#)

**Cost:** ~ \$3,000 per cycle

## Treatment schedule - Detail

The supportive therapies (e.g. antiemetics, premedications, etc.), infusion times, diluents, volumes and routes of administration, if included, are listed as defaults. They may vary between institutions and can be substituted to reflect individual institutional policy.

Antiemetics if included in the treatment schedule are based upon recommendations from national and international guidelines. These are **defaults only** and may be substituted to reflect individual institutional policy. Select here for recommended doses of alternative antiemetics.

## Cycle 1 and further cycles

Day 1		
Loratadine	10 mg (PO)	60 minutes before treatment
Dexamethasone	8 mg (PO)	60 minutes before treatment
Cetuximab	500 mg/m <sup>2</sup> (IV infusion)	over 2 hours (first dose); subsequent doses over 60 minutes *
Palonosetron	0.25 mg (IV bolus)	30 minutes before chemotherapy
Irinotecan	180 mg/m <sup>2</sup> (IV infusion)	in 250 mL to 500 mL glucose 5% over 90 minutes
Calcium folinate (Leucovorin)	50 mg (IV bolus)	over 1 to 2 minutes **
Fluorouracil	400 mg/m <sup>2</sup> (IV)	over 3 to 5 minutes
Fluorouracil	2,400 mg/m <sup>2</sup> (CIV)	via ambulatory infusion pump over 46 hours
Day 2 and 3		
Dexamethasone	8 mg (PO)	ONCE a day (or in divided doses) with or after food. Note: dexamethasone doses on day 2 and 3 may not be required and may be reduced or omitted at the clinicians discretion. ***

\* Although the product information recommends a maximum cetuximab infusion rate of 5 mg/min for the loading dose and 10 mg/min for subsequent doses, rates of 2 hours for the loading dose and 60 minutes for subsequent doses were used in clinical trials.<sup>1,2</sup>

\*\* The dose of calcium folinate (Leucovorin<sup>®</sup>) has been modified in this protocol from the original clinical trial dose of 200 mg/m<sup>2</sup> to 50 mg. A discussion regarding the effect of dosing on outcome can be found in the calcium folinate dose document.

\*\*\* Link to [ID 7 Prevention of chemotherapy induced nausea and vomiting](#)

**Frequency:** 14 days

**Cycles:** Continuous until disease progression or unacceptable toxicity

## Indications and patient population

- RAS wild-type metastatic colorectal cancer.

### Note:

- All patients should be tested for RAS mutations. Patients with mutant or unknown RAS status should not receive an EGFR antagonist as it may be harmful.
- Presence of a BRAF mutation has been identified as a marker of poorer prognosis, and potentially predictive of resistance to EGFR antagonists.<sup>3</sup>
- Consider side of primary tumour when prescribing treatment as patients with a right sided tumour may not benefit from the addition of an EGFR antagonist to chemotherapy.

## Clinical information

<b>Safety alert fluoropyrimidines</b>	<p>Fluoropyrimidines can be administered by different routes and schedules with each method having associated increased risk of certain side effects. Fluoropyrimidine overdose or overexposure is a rare but potentially life threatening side effect of this drug class and can occur by any route of administration. An antidote is available and highly effective if given within 96 hours.</p> <p>Read more about the <a href="#">medication safety alert for infusional fluorouracil and fluoropyrimidine overdose or overexposure</a></p>
<b>Venous access</b>	<p>Central venous access device (CVAD) is required to administer this treatment.</p> <p>Read more about <a href="#">central venous access device line selection</a></p>
<b>Hypersensitivity/infusion related reaction</b>	<p>High risk with cetuximab. The risk for anaphylactic reactions is increased in patients with a history of allergy to red meat or tick bites, or positive IgE antibody test results against cetuximab (<math>\alpha</math>-1-3- galactose).</p> <p>Read more about <a href="#">Hypersensitivity reaction</a></p>
<b>Premedication</b>	<p>The product information states that premedication is required for this treatment.</p> <p>Please refer to the treatment schedule for suggested premedication regimen. This may be substituted to reflect institutional policy.</p>
<b>Premedication</b>	<p>Consider atropine 0.3 to 0.6 mg IV or SC prior to irinotecan administration as required to help prevent irinotecan induced cholinergic side effects (atropine should not be used in patients with glaucoma).</p>
<b>Emetogenicity MODERATE</b>	<p>Suggested default antiemetics have been added to the treatment schedule, and may be substituted to reflect institutional policy.</p> <p>A steroid has been included both as an antiemetic and premedication for hypersensitivity in this protocol.</p> <p>For patients with a prior episode of chemotherapy induced nausea or vomiting, a NK1 receptor antagonist may be available on the PBS in combination with a 5HT<sub>3</sub> antagonist and steroid.</p> <p>Ensure that patients also have sufficient antiemetics for breakthrough emesis:</p> <p>Metoclopramide 10 mg three times a day when necessary (maximum of 30 mg/24 hours, up to 5 days) OR</p> <p>Prochlorperazine 10 mg PO every 6 hours when necessary.</p> <p>Read more about <a href="#">preventing anti-cancer therapy induced nausea and vomiting</a></p>
<b>Cardiac toxicity</b>	<p>Cardiac toxicity is a serious complication that can occur during treatment with fluorouracil. Patients treated with fluorouracil, especially those with a prior history of cardiac disease or other risk factors, should be carefully monitored during therapy.</p> <p>Read more about <a href="#">cardiac toxicity associated with anti-cancer drugs</a></p>
<b>Diarrhoea (early onset) and cholinergic syndrome</b>	<p>Early onset diarrhoea and other cholinergic symptoms such as rhinitis, hypersalivation, miosis, lacrimation, diaphoresis, flushing, and abdominal cramping can occur during or within 24 hours of administration of irinotecan.</p> <p>Symptoms may be treated with atropine 0.3 to 0.6 mg IV or SC as needed, repeated up to a maximum dose of 1.2 mg (unless clinically contraindicated). Patients 65 years of age and above should be closely monitored due to a greater risk of early diarrhoea observed in this population.</p> <p>Read more about <a href="#">treatment induced diarrhoea</a></p>
<b>Diarrhoea (late onset)</b>	<p>Irinotecan induced diarrhoea can be life threatening and requires immediate management.</p> <p>Ensure patients have sufficient antidiarrhoeal (e.g. loperamide) and appropriate instructions should this adverse event occur.</p> <p>Note: If prescribing loperamide, the recommended maximum daily dose of 16 mg of loperamide can be exceeded.</p> <p>Read more about <a href="#">treatment induced diarrhoea</a></p>

<b>Dihydropyrimidine dehydrogenase (DPD) enzyme deficiency</b>	<p>Rare, life-threatening toxicities such as mucositis, neutropenia, neurotoxicity and diarrhoea have been reported following administration of fluoropyrimidines (e.g. fluorouracil and capecitabine). Severe unexplained toxicities require investigation prior to continuing with treatment. Testing for DPD enzyme deficiency is available in Australia but not currently reimbursed.</p> <p>Read more about <a href="#">dihydropyrimidine dehydrogenase (DPD) enzyme deficiency</a></p>
<b>Severe enteropathy associated with fluoropyrimidine</b>	<p>Severe enteropathy has been reported among patients with stage II/III colon cancer treated with fluoropyrimidine chemotherapy with or without oxaliplatin. Patients treated with fluoropyrimidine should be closely monitored for diarrhoea and aggressively managed.</p> <p>Read more about <a href="#">severe enteropathy associated with fluorouracil in colorectal cancer</a></p>
<b>Gilbert's syndrome</b>	<p>Patients with Gilbert's syndrome should have their dose of irinotecan reduced. There is no clear dosing strategy; however based on the area under the concentration-time curve of SN-38, Innocenti et al (2006) recommend a 20% dose reduction of irinotecan.</p> <p>Read more about <a href="#">Gilbert's syndrome</a></p>
<b>Acneiform rash</b>	<p>EGFR targeted therapies are commonly associated with acneiform rash. The rash may peak in the first 2 to 4 weeks.</p> <p>Ensure advice on skin care (i.e. moisturisers) and sunscreen is provided. Prophylactic or early therapy with a tetracycline antibiotic (e.g. doxycycline) and 1% hydrocortisone cream to affected areas may be considered. Patients developing skin rash should be monitored for infectious sequelae, dose reductions and/or delay or cessation of treatment may be required.</p> <p>Read more about <a href="#">acneiform rash associated with EGFR inhibitors</a></p>
<b>Pulmonary toxicity</b>	<p>Interstitial lung disease (ILD) has been reported in patients treated with EGFR inhibitors.</p> <p>Read more about <a href="#">pulmonary toxicity associated with anti-cancer drugs</a>.</p>
<b>Blood tests</b>	<p>FBC, EUC, LFT's, calcium and magnesium at baseline and prior to each cycle. INR as clinically indicated. Magnesium wasting syndrome is associated with this therapy and patients should be monitored for hypomagnesaemia and accompanying hypocalcaemia for up to 8 weeks after completion of treatment.</p>
<b>Hepatitis B screening and prophylaxis</b>	<p>Routine screening for HBsAg and anti-HBc is NOT usually recommended for patients receiving this treatment.</p> <p>Read more about <a href="#">hepatitis B screening and prophylaxis in cancer patients requiring cytotoxic and/or immunosuppressive therapy</a></p>
<b>Vaccinations</b>	<p>Live vaccines are contraindicated in cancer patients receiving immunosuppressive therapy and/or who have poorly controlled malignant disease.</p> <p>Refer to the recommended schedule of vaccination for immunocompromised patients, as outlined in the <a href="#">Australian Immunisation Handbook</a>.</p> <p>Read more about <a href="#">COVID-19 vaccines and cancer</a>.</p>
<b>Fertility, pregnancy and lactation</b>	<p>Cancer treatment can have harmful effects on fertility and this should be discussed with all patients of reproductive potential prior to commencing treatment. There is a risk of foetal harm in pregnant women. A pregnancy test should be considered prior to initiating treatment in females of reproductive potential if sexually active. It is important that all patients of reproductive potential use effective contraception whilst on therapy and after treatment finishes. Effective contraception methods and adequate contraception timeframe should be discussed with all patients of reproductive potential. Possibility of infant risk should be discussed with breastfeeding patients.</p> <p>Read more about the <a href="#">effect of cancer treatment on fertility</a></p>

## Dose modifications

*Evidence for dose modifications is limited, and the recommendations made on eviQ are intended as a guide only. They are generally conservative with an emphasis on safety. Any dose modification should be based on clinical judgement, and the individual patient's situation including but not limited to treatment intent (curative vs palliative), the anti-cancer regimen (single*

versus combination therapy versus chemotherapy versus immunotherapy), biology of the cancer (site, size, mutations, metastases), other treatment related side effects, additional co-morbidities, performance status and patient preferences. Suggested dose modifications are based on clinical trial findings, product information, published guidelines and reference committee consensus. The dose reduction applies to each individual dose and not to the total number of days or duration of treatment cycle unless stated otherwise. Non-haematological gradings are based on [Common Terminology Criteria for Adverse Events \(CTCAE\)](#) unless otherwise specified. Renal and hepatic dose modifications have been standardised where possible. For more information see dosing considerations & disclaimer.

The dose recommendations in kidney dysfunction (i.e. renal impairment) displayed may not reflect those in the ADDIKD guideline and have been included for historical reference only. Recommendations will be updated once the individual protocol has been evaluated by the reference committee, with this version of the protocol then being archived. Clinicians are expected to refer to the ADDIKD guideline prior to prescribing in kidney dysfunction.

[International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction \(ADDIKD\)](#).

**Note:**

- All dose reductions are calculated as a percentage of the starting dose.
- The dose of calcium folinate (Leucovorin®) remains fixed at 50 mg and is delayed or omitted if fluorouracil bolus is delayed or omitted.

Haematological toxicity	
ANC x 10 <sup>9</sup> /L (pre-treatment blood test)	
1.0 to less than 1.5	Refer to local institutional guidelines; it is the view of the expert clinicians that treatment should continue if patient is clinically well.
0.5 to less than 1.0	Delay treatment until recovery
less than 0.5	Delay treatment until recovery and consider reducing irinotecan and fluorouracil by 25% for subsequent cycles
Febrile neutropenia	Delay treatment until recovery and consider reducing irinotecan and fluorouracil by 25% for subsequent cycles
Platelets x 10 <sup>9</sup> /L (pre-treatment blood test)	
75 to less than 100	The general recommendation is to delay, however if the patient is clinically well it may be appropriate to continue treatment; refer to treating team and/or local institutional guidelines.
50 to less than 75	Delay treatment until recovery
less than 50	Delay treatment until recovery and consider reducing irinotecan and fluorouracil by 25% for subsequent cycles

Renal impairment	
Creatinine clearance (mL/min)	
30 to 50	Reduce irinotecan and fluorouracil by 25%
less than 30	Reduce irinotecan and fluorouracil by 50%

Hepatic impairment	
Hepatic dysfunction	
Minimal	Reduce irinotecan by 25%
Mild	Reduce irinotecan by 50%
Moderate	Omit irinotecan and reduce fluorouracil by 25% or withhold chemotherapy
Severe	Omit irinotecan and reduce fluorouracil by 50% or withhold chemotherapy
<a href="#">Gilbert's syndrome</a>	Reduce irinotecan by 20%

**Mucositis and stomatitis**

Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce fluorouracil by 25% 3 <sup>rd</sup> occurrence: Reduce fluorouracil by 50% 4 <sup>th</sup> occurrence: Omit fluorouracil
Grade 3 or Grade 4	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce fluorouracil by 50% 2 <sup>nd</sup> occurrence: Omit fluorouracil

### Diarrhoea

Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce doses for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce irinotecan, fluorouracil and cetuximab by 25% 3 <sup>rd</sup> occurrence: Reduce irinotecan, fluorouracil and cetuximab by 50% 4 <sup>th</sup> occurrence: Cease chemotherapy
Grade 3 or Grade 4	Delay treatment until toxicity has resolved to Grade 1 or less and reduce doses for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce irinotecan and fluorouracil by 50% 2 <sup>nd</sup> occurrence: Cease chemotherapy

### Hand foot syndrome (Palmar-plantar erythrodysesthesia)

Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce fluorouracil 25% 3 <sup>rd</sup> occurrence: Reduce fluorouracil by 50% 4 <sup>th</sup> occurrence: Omit fluorouracil
Grade 3	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce fluorouracil by 50% 2 <sup>nd</sup> occurrence: Omit fluorouracil

### Rash acneiform

Grade 3	Delay treatment until toxicity has resolved to Grade 2 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce cetuximab by 25% 3 <sup>rd</sup> occurrence: Reduce cetuximab by 50% 4 <sup>th</sup> occurrence: Omit cetuximab
---------	---

## Interactions

Drug interactions in eviQ protocols are under review and being updated to align with current literature. Further site-wide updates and changes will occur in due course. *References & Disclaimer*

The drug interactions shown below are not an exhaustive list. For a more comprehensive list and for detailed information on specific drug interactions and clinical management, please refer to the specific drug product information and the following key resources:

- [MIMS - interactions tab](#) (includes link to a CYP-450 table) (login required)
- [Australian Medicines Handbook \(AMH\) – interactions tab](#) (login required)
- [Micromedex Drug Interactions](#) (login required)
- [Cancer Drug Interactions](#)

<b>Cetuximab</b>		
	<b>Interaction</b>	<b>Clinical management</b>
<b>Chemotherapeutic agents</b>	Increased incidence of specific adverse reactions when used in combination	Monitor closely (e.g. for cardiac toxicity and hand-foot syndrome when combined with fluoropyrimidines; for severe diarrhoea with capecitabine and oxaliplatin)

<b>Fluorouracil</b>		
	<b>Interaction</b>	<b>Clinical management</b>
<b>Folic acid</b>	Increased toxicity of fluorouracil due to stabilisation of its bond to thymidylate synthetase (folic acid is a precursor of folinic acid/leucovorin)	Advise patients not to take folic acid supplements (inc. multivitamins) around the time of receiving treatment with fluorouracil
<b>Metronidazole, tinidazole</b>	Increased toxicity of fluorouracil due to reduced clearance	Avoid combination or monitor for fluorouracil toxicity
<b>Warfarin and other drugs metabolised by CYP2C9 (e.g. warfarin, phenytoin etc.)</b>	Increased effect/toxicity of these drugs due to inhibition of CYP2C9 by fluorouracil resulting in reduced clearance	Avoid combination or monitor for increased effect/toxicity of these drugs (e.g. for bleeding/elevated INR with warfarin, elevated phenytoin serum levels or signs of toxicity such as ataxia, tremor etc.)
<b>Allopurinol</b>	Reduced efficacy of fluorouracil possible due to reduced conversion to the active metabolites	Avoid combination or monitor for reduced fluorouracil efficacy

<b>Irinotecan</b>		
	<b>Interaction</b>	<b>Clinical management</b>
<b>CYP3A4 and P-gp inhibitors (e.g. amiodarone, aprepitant, azole-antifungals, ritonavir, lapatinib, nilotinib, sorafenib, macrolides, ciclosporin, grapefruit juice etc.)</b>	Increased toxicity of irinotecan possible due to reduced clearance	Avoid combination or monitor for irinotecan toxicity  (Ketoconazole contraindicated and should be discontinued at least 1 week prior to irinotecan)
<b>CYP3A4 and P-gp inducers (e.g. carbamazepine, phenytoin, phenobarbitone, rifampicin, St John's wort, dexamethasone etc.)</b>	Reduced efficacy of irinotecan possible due to increased clearance	Avoid combination or monitor for decreased clinical response to irinotecan  (St John's wort contraindicated; substituting non-enzyme inducing antiepileptics (e.g. clonazepam, diazepam, lorazepam) at least 1 week prior to irinotecan should be considered)
<b>Atazanavir</b>	Increased toxicity of irinotecan possible due to reduced clearance caused by inhibition of both CYP3A4 and UGT1A1 by atazanavir	Avoid combination or monitor for irinotecan toxicity
<b>Smoking</b>	Reduced efficacy of irinotecan possible due to increased clearance caused by induction of both CYP3A4 and UGT1A1 by smoking	Monitor for decreased clinical response to irinotecan in patients who continue to smoke; no specific dosing recommendations are available

General		
	Interaction	Clinical management
<b>Warfarin</b>	Anti-cancer drugs may alter the anticoagulant effect of warfarin.	Monitor INR regularly and adjust warfarin dosage as appropriate; consider alternative anticoagulant.
<b>Direct oral anticoagulants (DOACs) e.g. apixaban, rivaroxaban, dabigatran</b>	Interaction with both CYP3A4 and P-gp inhibitors /inducers.  DOAC and anti-cancer drug levels may both be altered, possibly leading to loss of efficacy or toxicity (i.e. increased bleeding).	Apixaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. If treating VTE, avoid use with strong CYP3A4 and P-gp inducers.  Rivaroxaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors.  Dabigatran: avoid combination with strong P-gp inducers and inhibitors.  If concurrent use is unavoidable, monitor closely for efficacy/toxicity of both drugs.
<b>Digoxin</b>	Anti-cancer drugs can damage the lining of the intestine; affecting the absorption of digoxin.	Monitor digoxin serum levels; adjust digoxin dosage as appropriate.
<b>Antiepileptics</b>	Both altered antiepileptic and anti-cancer drug levels may occur, possibly leading to loss of efficacy or toxicity.	Where concurrent use of an enzyme-inducing antiepileptic cannot be avoided, monitor antiepileptic serum levels for toxicity, as well as seizure frequency for efficacy; adjust dosage as appropriate. Also monitor closely for efficacy of the anti-cancer therapy.
<b>Antiplatelet agents and NSAIDs</b>	Increased risk of bleeding due to treatment related thrombocytopenia.	Avoid or minimise combination. If combination deemed essential, (e.g. low dose aspirin for ischaemic heart disease) monitor for signs of bleeding.
<b>Serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs e.g. paroxetine) and serotonin noradrenaline reuptake inhibitors (SNRIs e.g. venlafaxine)</b>	Increased risk of serotonin syndrome with concurrent use of 5-HT3 receptor antagonists (e.g. palonosetron, ondansetron, granisetron, tropisetron, dolasetron, etc.)	Avoid combination. If combination is clinically warranted, monitor for signs and symptoms of serotonin syndrome (e.g. confusion, agitation, tachycardia, hyperreflexia). For more information link to <a href="#">TGA Medicines Safety Update</a>
<b>Vaccines</b>	Diminished response to vaccines and increased risk of infection with live vaccines.	Live vaccines (e.g. BCG, MMR, zoster and varicella) are contraindicated in patients on immunosuppressive therapy. Use with caution in patients on non-immunosuppressive therapy. For more information; refer to the recommended schedule of vaccination for cancer patients, as outlined in the <a href="#">Australian Immunisation Handbook</a>

## Administration

*eviQ provides safe and effective instructions on how to administer cancer treatments. However, eviQ does not provide every treatment delivery option, and is unable to provide a comprehensive list of cancer treatment agents and their required IV line giving set/filter. There may be alternative methods of treatment administration, and alternative supportive treatments that are also appropriate. Please refer to the individual*



## Day 1

### Approximate treatment time: 4 to 5 hours

#### Safe handling and waste management

#### Safe administration

General patient assessment prior to each treatment.

Any toxicity grade 2 or greater may require dose reduction, delay or omission of treatment and review by medical officer before recommencing treatment.

Prime IV line(s).

Access TIVAD or CVAD.

### Pre treatment medication

Verify premedication taken or administer as prescribed.

Verify dexamethasone taken or administer as prescribed.

## 🕒 Treatment - Time out

### Cetuximab

- administer via IV infusion over 2 hours (first dose)
- if well tolerated, subsequent doses may be administered over 60 minutes
- although the product information recommends a maximum cetuximab infusion rate of 5 mg/min for the loading dose and 10 mg/min for subsequent doses, rates of 2 hours for the loading dose and 60 minutes for subsequent doses were used in clinical trials
- observe for hypersensitivity reaction
- flush with ~ 50 mL of sodium chloride 0.9%
- patient should be observed for an hour post infusion. If patient has a hypersensitivity reaction – stop infusion immediately. Review by medical officer; if re-challenge indicated, pre medicate patient and recommence infusion over 2 hours , then patient should be observed for an hour post infusion.

*The product information recommends waiting one hour post completion of the cetuximab infusion, before commencing concomitant chemotherapy. However, to reflect clinical practice, if no reactions are observed with the first two doses, concomitant chemotherapy may be administered immediately following cetuximab at the discretion of the treating physician.*

### Pre treatment medication

Verify antiemetics taken or administer as prescribed.

## 🕒 Chemotherapy - Time out

### Irinotecan

#### Prior to administration:

- administer atropine premedication if required
- patient vital signs should be closely monitored post atropine administration as atropine can cause tachycardia, heart arrhythmias, hypertension and angina.

#### Administer irinotecan (irritant):

- via IV infusion over 90 minutes
- protect from light
- flush with ~ 100 mL of sodium chloride 0.9%
- observe patient for cholinergic symptoms
- if patient develops early onset diarrhoea and other cholinergic symptoms (such as rhinitis, increased salivation, miosis, lacrimation, diaphoresis, flushing) - stop infusion, monitor vital signs and review by medical officer.

## Calcium Folate (Leucovorin)

- administer by IV bolus via a side port of the IV line over 1 to 2 minutes
- flush with ~ 50mL of sodium chloride 0.9%.

## Fluorouracil

- If using cryotherapy commence ~ 5 minutes prior to administering fluorouracil and continue for ~ 30 minutes post.

### Administer fluorouracil (irritant):

- over 3 to 5 minutes
  - via a minibag **OR**
  - by IV bolus via a side port of a freely flowing IV infusion
- flush with ~ 100 mL of sodium chloride 0.9%.

## Fluorouracil continuous infusion (irritant)

Connect pump containing fluorouracil and administer over the correct time for the amount of drug in the pump:

- **▲ safety alert** issued for administration of infusional fluorouracil
- verify the correct rate of infusion via the ambulatory infusion pump
- read more information about the different [ambulatory infusion pumps](#).

Continue **safe handling** precautions until 7 days after completion of drug(s)

---

## Day 3

Approximate treatment time: 30 minutes

[Safe handling and waste management](#)

### Disconnection of ambulatory infusion pump/infusor

Verify the ambulatory infusion pump/infusor is complete.

Disconnect the ambulatory infusion pump/infusor as per recommended procedure for type of pump/infusor.

Read more about [ambulatory infusion pumps/infusors](#).

Deaccess [TIVAD](#) or [CVAD](#).

Continue **safe handling** precautions until 7 days after completion of drug(s)

---

## Discharge information

### Antiemetics

- Antiemetics as prescribed.

### Antidiarrhoeals

- Antidiarrhoeals (e.g. loperamide) as prescribed with written instructions on how to manage this side effect and 24 hour emergency contact.

### Patient information

- Ensure patient receives patient information sheet.

### Infusion pumps

- CADD-Legacy® 1 ambulatory infusion pump [patient information sheet](#).
- CADD-Legacy® Plus ambulatory infusion pump [patient information sheet](#).
- CADD® Solis VIP ambulatory infusion pump [patient information sheet](#).
- Elastomeric infusion system [patient information sheet](#).

## Side effects

*The side effects listed below are not a complete list of all possible side effects for this treatment. Side effects are categorised into the*

Immediate (onset hours to days)	
<b>Hypersensitivity reaction</b>	Anaphylaxis and infusion related reactions can occur with this treatment. Read more about <a href="#">hypersensitivity reaction</a>
<b>Diarrhoea and cholinergic syndrome (early onset) associated with irinotecan</b>	Cholinergic syndrome can occur during or shortly after commencing the irinotecan infusion, or within 24 hours of administration of the drug. It is characterised by diarrhoea, rhinitis, hypersalivation, miosis, lacrimation, diaphoresis, flushing, and abdominal cramping. Read more about <a href="#">treatment induced diarrhoea</a>
<b>Nausea and vomiting</b>	Read more about <a href="#">prevention of treatment induced nausea and vomiting</a>
<b>Taste and smell alteration</b>	Read more about <a href="#">taste and smell changes</a>
<b>Cardiotoxicity</b>	Coronary artery spasm is a temporary, sudden narrowing of one of the coronary arteries that may present at any time during treatment with fluoropyrimidines. It most commonly manifests as angina.
Early (onset days to weeks)	
<b>Neutropenia</b>	Abnormally low levels of neutrophils in the blood. This increases the risk of infection. Any fever or suspicion of infection should be investigated immediately and managed aggressively. Read more about <a href="#">immediate management of neutropenic fever</a>
<b>Thrombocytopenia</b>	A reduction in the normal levels of functional platelets, increasing the risk of abnormal bleeding.  Read more about <a href="#">thrombocytopenia</a>
<b>Diarrhoea (late onset) associated with irinotecan</b>	Late onset of diarrhoea after 24 hours post irinotecan administration can be life threatening and requires immediate treatment. Read more about <a href="#">treatment induced diarrhoea</a>
<b>Anorexia</b>	Loss of appetite accompanied by decreased food intake. Read more about <a href="#">anorexia</a>
<b>Oral mucositis</b>	Erythematous and ulcerative lesions of the gastrointestinal tract (GIT). It commonly develops following chemotherapy, radiation therapy to the head, neck or oesophagus, and high dose chemotherapy followed by a blood and marrow transplant (BMT). Read more about <a href="#">oral mucositis</a>
<b>Fatigue</b>	Read more about <a href="#">fatigue</a>
<b>Ocular changes</b>	Symptoms may include eye pain, blurred vision, blepharitis, uveitis, optic neuritis, tear duct stenosis, conjunctivitis, hyperlacrimation, watery or dry eyes and photophobia.
<b>Acneiform rash</b>	A skin rash, characterised by papules and pustules affecting the face and upper body. This is commonly associated with small molecule EGFR inhibitors and some monoclonal antibodies (e.g. cetuximab, panitumumab). Read more about <a href="#">acneiform rash associated with EGFR inhibitors</a>
<b>Actinic keratoses flare</b>	Pre-existing actinic keratoses (AKs) can become more inflamed and scaly as a result of immunosuppression. Read more about <a href="#">actinic keratoses flare</a>
<b>Photosensitivity</b>	Increased sensitivity to ultraviolet (UV) light resulting in an exaggerated sunburn-like reaction accompanied by stinging sensations and urticaria.
<b>Palmar-plantar erythrodysesthesia (PPE) - hand-foot syndrome (HFS)</b>	Bilateral erythema, tenderness, pain, swelling, tingling, numbness, pruritus, dry rash, or moist desquamation and ulceration of the palms and soles. It is also known as hand-foot syndrome (HFS). Symptoms appear to be dose dependent and palms are affected more than soles. Read more about <a href="#">hand-foot syndrome associated with chemotherapy</a>

Late (onset weeks to months)	
<b>Anaemia</b>	Abnormally low levels of red blood cells (RBCs) or haemoglobin in the blood. Read more about <a href="#">anaemia</a>
<b>Paronychia</b>	An inflammatory reaction involving the folds of the skin surrounding the fingernail. Read about <a href="#">nail toxicities</a>
<b>Abnormal hair growth</b>	Hair may become fine, brittle and curly. Eyelashes and eyebrows may grow more quickly and become unusually long.
<b>Hyperpigmentation</b>	Darkening of an area of skin caused by the overproduction of melanin.
<b>Pulmonary toxicity</b>	Pulmonary toxicity may include damage to the lungs, airways, pleura and pulmonary circulation. Read more about <a href="#">pulmonary toxicity associated with anti-cancer drugs</a>

## Evidence

Due to the lack of conclusive evidence to identify the optimum dose of calcium folinate (Leucovorin®), it is the consensus of the eviQ reference committee to adopt flat dosing of calcium folinate (Leucovorin®) as a 50 mg IV bolus when used with bolus 5FU across all colorectal and upper gastrointestinal protocols. A discussion regarding the effect of dosing on outcome can be found in the calcium folinate (Leucovorin®) dose document.

The evidence supporting the use of FOLFIRI and cetuximab in the first-line treatment for metastatic colorectal cancer is based on the CRYSTAL study, a randomized, open-label, multicentre study that compared cetuximab plus FOLFIRI and FOLFIRI alone in 1,198 patients. The patients receiving cetuximab plus FOLFIRI received cetuximab at 400 mg/m<sup>2</sup> on day 1 and subsequently received cetuximab at 250 mg/m<sup>2</sup> weekly. This was given together with FOLFIRI on a 14 day cycle.<sup>4</sup>

Monoclonal antibodies (mABs) targeting the epidermal growth factor receptor (EGFR) prolong survival in patients with metastatic colorectal cancer (mCRC) harbouring KRAS exon 2 wild type tumours. Recent evidence suggest that other RAS mutations (exon 3 and 4 of KRAS and exons 2, 3, 4 of NRAS) may also be predictive of resistance<sup>5,6</sup> and some studies even suggest that anti-EGFR mAB treatment may have a detrimental effect on PFS and OS.<sup>7</sup> As such, cetuximab should not be used in patients with any RAS mutations.

Several studies have shown that the 2 weekly schedule of cetuximab has similar efficacy and safety as the weekly schedule and is sometimes preferred as it is more convenient for the patient.<sup>8</sup> However, at present there are no published phase III studies to confirm this.

In the controlled phase 2 APEC study, it was shown that there was similar efficacy and safety profile for 2 weekly cetuximab compared with historical data on weekly schedules.<sup>9</sup>

In the CECOG randomised multicentre phase II study the combination of cetuximab every second week with FOLFOX4 demonstrated activity and a manageable safety profile comparable with that observed for FOLFOX4 plus standard weekly cetuximab.<sup>10</sup>

Another phase 2 study by Martin-Martorelli et al also showed similar efficacy for cetuximab every two weeks in combination with irinotecan.<sup>1</sup>

In the second part of the Phase 1 dose escalation study by Tabernero et al, cetuximab two weekly was administered in combination with FOLFIRI. The primary end-point of this study was to determine the maximum tolerated dose of cetuximab administered two weekly, however efficacy data was reported and did not show any significant differences to the weekly regimen.<sup>2</sup>

The expert reference panel supported publication of the protocol on the basis of the information summarised below.

Source	Study & Year Published	Supports Use	Is the dose and regimen consistent with the protocol?	Comments
Phase II trials	Cheng et al 2013 <sup>9</sup>	Yes	Yes	-
	Martin-Martorelli et al 2008 <sup>1</sup>	Yes	No	Irinotecan/Cetuximab
	Brodowicz et al	Yes	No	FOLFOX + cetuximab

Source	Study & Year Published	Supports Use	Is the dose and regimen consistent with the protocol?	Comments
	2013 <sup>10</sup>			weekly vs FOLFOX + cetuximab 2 weekly
Phase I trial	Taberero et al 2010 <sup>2</sup>	Yes	Yes	Population pK/pD dose escalation study
Observational studies	-	N/A	-	-
Case series	Pfeiffer et al 2008 <sup>11</sup>	Yes	No	Irinotecan/Cetuximab
Guidelines	Date published/revised	Supports Use	Is the dose and regimen consistent with the protocol?	Comments
NCCN	v2. 2015	Yes	Yes	-
BCCA	-	N/A	-	-
CCO	-	N/A	-	-

## Efficacy



In a small phase 1 dose escalation study of cetuximab monotherapy using bi-weekly dosing, efficacy data was looked at and there were comparable results for cetuximab every two weeks.<sup>2</sup>

### Cetuximab weekly/FOLFIRI

After a median follow up of 46.8 months, the addition of cetuximab to FOLFIRI resulted in an improvement in overall survival time from 18.6 months to 19.9 months (HR 0.878, 95% CI, 0.774 to 0.995; P = 0.0419).<sup>12</sup>

A post-hoc analysis investigating the impact of RAS mutations other than KRAS codon 12 or 13 in relation to treatment effects showed that there was no clear evidence that the addition of cetuximab to FOLFIRI modified efficacy outcome in the evaluable patients with other tumour RAS mutations. In patients with RAS wild-type tumors, a clear cetuximab benefit was seen across efficacy end points.<sup>5</sup>

**Kaplan-Meier plots for overall survival according to treatment group in RAS populations. (A) KRAS codon 12 or 13 wild type, evaluable for other RAS mutations. (B) RAS wild type (all loci). (C) KRAS codon 12 or 13 wild type; other RAS mutations. (D) RAS mutation (any locus)<sup>5</sup>**





No. at risk

	0	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57
CT + cetuximab	210	202	187	176	162	158	145	121	107	96	79	70	60	56	48	32	17	5	3	0
CT	220	212	196	183	156	133	114	97	88	76	62	51	41	34	30	21	11	6	2	0

No. at risk

	0	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57
CT + cetuximab	178	174	163	155	142	140	128	108	97	89	73	66	56	52	45	29	16	5	3	0
CT	189	182	171	160	135	115	98	85	79	70	58	47	38	32	28	20	10	6	2	0



No. at risk


	0	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51
CT + cetuximab	32	28	24	21	20	18	17	13	10	7	6	4	4	4	3	3	1	0
CT	31	30	25	23	21	18	16	12	9	6	4	4	3	2	2	1	1	0

No. at risk

	0	3	6	9	12	15	18	21	24	27	30	33	36	39	42	45	48	51	54	57
CT + cetuximab	246	223	205	186	164	134	107	84	66	55	43	38	36	32	27	22	12	3	2	1
CT	214	205	189	168	148	113	93	72	59	51	42	38	32	23	20	13	3	1	1	0

@ Journal of Clinical Oncology 2015

Hazard ratios for (A) overall survival and (B) progression-free survival according to tumour KRAS exon 2 and RAS mutation status



@ Journal of Clinical Oncology 2015

## Toxicity

In the studies looking at the biweekly cetuximab combination regimens, there were no increased adverse events, including allergic reactions and skin toxicity associated with the double dose.<sup>8</sup>

## Toxicity of Cetuximab (Weekly) plus FOLFIRI vs FOLFIRI Alone<sup>4</sup>

**Table 3. Most Common Grade 3 or 4 Adverse Events and Special Adverse Event Categories in the Safety Population, According to Treatment Group.<sup>2</sup>**

MedDRA Preferred Term <sup>†</sup>	Cetuximab plus FOLFIRI (N = 600) no. (%)	FOLFIRI Alone (N = 602) no. (%)	P Value
Any	476 (79.3)	367 (61.0)	<0.001
Neutropenia <sup>‡</sup>	169 (28.2)	148 (24.6)	0.16
Leukopenia	43 (7.2)	31 (5.1)	0.15
Diarrhea	94 (15.7)	63 (10.5)	0.008
Fatigue	32 (5.3)	28 (4.7)	0.59
Rash	49 (8.2)	0	<0.001
Dermatitis acneiform	32 (5.3)	0	<0.001
Vomiting	28 (4.7)	30 (5.0)	0.80
Special adverse events			
Skin reactions:			
All	118 (19.7)	1 (0.2)	<0.001
Acne-like rash	97 (16.2)	0	<0.001
Infusion-related reaction	15 (2.5)	0	<0.001

\* We used retrospective chi-square tests to compare the rates of adverse events between the two treatment groups; the results were not corrected for multiple testing. Under the assumption of no significant difference between the two groups, 11 independent tests and a 0.05 significance level results in a chance of more than 43% of obtaining at least one false positive finding. FOLFIRI denotes irinotecan, fluorouracil, and leucovorin.  
<sup>†</sup> Among the *Medical Dictionary for Regulatory Activities* (MedDRA, version 10.0) preferred terms, no grade 4 reactions were reported for dermatitis acneiform, acne-like rash, or all skin reactions.  
<sup>‡</sup> Grade 3 or 4 febrile neutropenia was reported in 18 of the 600 patients (3.0%) receiving cetuximab plus FOLFIRI and in 13 of the 602 patients (2.2%) receiving FOLFIRI alone.

© New England Journal of Medicine 2009

## References

- Martin-Martorell, P., S. Rosello, E. Rodriguez-Braun, et al. 2008. "Biweekly cetuximab and irinotecan in advanced colorectal cancer patients progressing after at least one previous line of chemotherapy: results of a phase II single institution trial." *Br J Cancer* 99(3):455-458.
- Taberero, J., F. Ciardiello, F. Rivera, et al. 2010. "Cetuximab administered once every second week to patients with metastatic colorectal cancer: a two-part pharmacokinetic/pharmacodynamic phase I dose-escalation study." *Ann Oncol* 21(7):1537-1545.
- Nott, L., M. Khattak, T. Price, et al. Molecular pathology and biomarkers – implications for systemic therapy. Available from [https://wiki.cancer.org.au/australia/Guidelines:Colorectal\\_cancer/Systemic\\_therapy\\_molecular\\_pathology](https://wiki.cancer.org.au/australia/Guidelines:Colorectal_cancer/Systemic_therapy_molecular_pathology). In: Cancer Council Australia Colorectal Cancer Guidelines Working Party.
- Van Cutsem, E., C. H. Kohne, E. Hitre, et al. 2009. "Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer." *N Engl J Med* 360(14):1408-1417.
- Van Cutsem, E., H. J. Lenz, C. H. Kohne, et al. 2015. "Fluorouracil, Leucovorin, and Irinotecan Plus Cetuximab Treatment and RAS Mutations in Colorectal Cancer." *J Clin Oncol* 33(7):692-700.
- Heinemann, V., L. F. von Weikersthal, T. Decker, et al. 2014. "FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial." *Lancet Oncol* 15(10):1065-1075.
- Douillard, J. Y., K. S. Oliner, S. Siena, et al. 2013. "Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer." *N Engl J Med* 369(11):1023-1034.
- Hubbard, J. M. and S. R. Alberts. 2013. "Alternate dosing of cetuximab for patients with metastatic colorectal cancer." *Gastrointest Cancer Res* 6(2):47-55.
- Cheng, Ann-Lii, Gerardo Cornelio, Lin Shen, et al. 2013. "First-line Cetuximab with FOLFOX or FOLFIRI Every 2 Weeks In KRAS Wild-Type Metastatic Colorectal Cancer: phase II APEC Study." *Annals of oncology* 24 (suppl 4):iv34-iv35. Abstract No:PD-0028
- Brodowicz, T., T. E. Ciuleanu, D. Radosavljevic, et al. 2013. "FOLFOX4 plus cetuximab administered weekly or every second

week in the first-line treatment of patients with KRAS wild-type metastatic colorectal cancer: a randomized phase II CECOG study." *Ann Oncol* 24(7):1769-1777.

- 11 Pfeiffer, P., D. Nielsen, J. Bjerregaard, et al. 2008. "Biweekly cetuximab and irinotecan as third-line therapy in patients with advanced colorectal cancer after failure to irinotecan, oxaliplatin and 5-fluorouracil." *Ann Oncol* 19(6):1141-1145.
- 12 Van Cutsem, E., C. H. Kohne, I. Lang, et al. 2011. "Cetuximab plus irinotecan, fluorouracil, and leucovorin as first-line treatment for metastatic colorectal cancer: updated analysis of overall survival according to tumor KRAS and BRAF mutation status." *J Clin Oncol* 29(15):2011-2019.

## History

### Version 5

Date	Summary of changes
24/03/2021	Cetuximab infusion rate information in detailed treatment schedule and administration section updated to include maximum infusion rates as per product information. Version number increased to V.5.
28/07/2021	Protocol title updated to "Colorectal metastatic FOLFIRI (modified) (fluorouracil leucovorin irinotecan) and cetuximab (two weekly)".
20/10/2022	Protocol reviewed electronically by Medical Oncology Reference Committee. No changes. Next review 2 years.

### Version 4

Date	Summary of changes
04/12/2020	Cetuximab hypersensitivity/infusion related reaction clinical information updated to include risk factors as per Medical Oncology Reference Committee meeting 23 <sup>rd</sup> October 2020. Version number increased to V.4. Next review in 2 years.

### Version 3

Date	Summary of changes
27/03/2015	New protocol taken to Medical Oncology Reference Committee meeting.
27/04/2015	Approved and published on eviQ.
18/02/2016	Discussion with Medical Oncology Reference Committee Chairs and protocol to be reviewed every 5 years. Next review due in 4 years.
16/10/2016	Patient information sheet updated to include more fluorouracil toxicity symptom warnings.
09/11/2016	The following changes made post Medical Oncology Reference Committee meeting held on 21 October 2016. Link to AGTIG and ANZCTR added. ANC dose modifications changed to 0.5 to less than 1.5 and less than 0.5. Dose reduction for ANC less than 0.5 and febrile neutropenia changed to 25%. Platelets dose modifications changed to 50 to less than 100 and less than 50. Sentence in dose modifications regarding omitting leucovorin if fluorouracil is delayed or omitted changed to specify fluorouracil bolus.
19/12/2016	The following sentence added to Indications and Patient population after discussion at Medical Oncology Reference Committee meeting held on 21 October 2016: Consider BRAF mutation status and side of primary tumour when prescribing treatment as patients with a BRAF mutation and/or right sided tumour may not benefit from the addition of cetuximab to chemotherapy and another regimen should be considered.
31/05/2017	Transferred to new eviQ website. Version number change to V.2.  Hepatitis screening changed to not recommended.
07/03/2018	Fluoropyrimidine warning added. Note in indication on BRAF updated by Medical Oncology Reference Committee.
10/05/2018	Haematological dose modifications updated as per consensus of the expert clinician group.



Date	Summary of changes
	Fluoropyrimidine and DPD enzyme deficiency wording in clinical information updated. Version number changed to V.3.
22/06/2018	Antiemetics updated to be in line with international guidelines. Note to dexamethasone added.
04/07/2018	Indications reworded for consistency across all colorectal EGFR monoclonal antibody protocols.
25/09/2018	Treatment schedule note and evidence section updated with Leucovorin <sup>®</sup> dosing information as per reference committee consensus.
25/09/2020	Protocol reviewed electronically by the Medical Oncology Reference committee. Nil changes. Next review in 2 years.

The information contained in this protocol is based on the highest level of available evidence and consensus of the eviQ reference committee regarding their views of currently accepted approaches to treatment. Any clinician (medical oncologist, haematologist, radiation oncologist, medical physicist, radiation therapist, pharmacist or nurse) seeking to apply or consult this protocol is expected to use independent clinical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use is subject to eviQ's disclaimer available at [www.eviq.org.au](http://www.eviq.org.au)

**First approved:** 27 April 2015  
**Last reviewed:** 20 October 2022  
**Review due:** 31 December 2024

***The currency of this information is guaranteed only up until the date of printing, for any updates please check:***

<https://www.eviq.org.au/p/1682>  
12 Jul 2023

# Patient information - Bowel cancer metastatic - FOLFIRI modified (fluorouracil, leucovorin, irinotecan) and cetuximab (two weekly)

Patient's name:


## Your treatment

The treatment schedule below explains how the drugs for this treatment are given.

FOLFIRI modified (fluorouracil, leucovorin, irinotecan) and cetuximab			
This treatment cycle is repeated every 14 days. Your doctor will advise you of the number of treatments you will have.			
Day	Treatment	How it is given	How long it takes
1	<b>Cetuximab</b> (se-TUK-see-mab)  <b>Irinotecan</b> (eye-ri-noe-TEE-kan)  <b>Calcium folinate (Leucovorin)</b> (loo-koe-VOR-in)  <b>Fluorouracil</b> (Flure-oh-YOOR-a-sill)	By a drip into a vein	About 4 to 5 hours
	<b>Fluorouracil</b> (Flure-oh-YOOR-a-sill)	By a pump slowly into a vein	For 2 days (46 hours) by pump at home
3	<b>Disconnect pump</b>		About 30 minutes

## When to get help

Anticancer drugs (drugs used to treat cancer) can sometimes cause serious problems. It is important to get medical help immediately if you become unwell.

	<p><b>IMMEDIATELY go to your nearest hospital Emergency Department, or contact your doctor or nurse if you have any of the following at any time:</b></p>	<p><b>Emergency contact details</b></p> <p>Ask your doctor or nurse from your treating team who to contact if you have a problem</p>
<ul style="list-style-type: none"> <li>• a temperature of 38°C or higher</li> <li>• chills, sweats, shivers or shakes</li> <li>• shortness of breath</li> <li>• uncontrolled vomiting or diarrhoea</li> <li>• pain, tingling or discomfort in your chest or arms</li> <li>• leaking from your pump</li> <li>• you become unwell.</li> </ul>	<p>Daytime:.....</p> <p>Night/weekend:.....</p> <p>Other instructions:.....</p> <p>.....</p> <p>.....</p> <p>.....</p>	

**During your treatment immediately** tell the doctor or nurse looking after you if you get any of the following problems:

- leaking from the area where the drugs are being given
- pain, stinging, swelling or redness in the area where the drugs are being given or at any injection sites
- a skin rash, itching, feeling short of breath, wheezing, fever, shivers, or feeling dizzy or unwell in any way (allergic reaction).

## Other information about your treatment

### Changes to your dose or treatment delays

Sometimes a treatment may be started at a lower dose or the dose needs to be changed during treatment. There may also be times when your treatment is delayed. This can happen if your doctor thinks you are likely to have severe side effects, if you get severe side effects, if your blood counts are affected and causing delays in treatment, or if you are finding it hard to cope with the treatment. This is called a dose reduction, dose change or treatment delay. Your doctor will explain if you need any changes or delays to your treatment and the reason why.

### Blood tests and monitoring

Anti-cancer drugs can reduce the number of blood cells in your body. You will need to have regular blood tests to check that your blood cell count has returned to normal. If your blood count is low your treatment may be delayed until it has returned to normal. Your doctor or nurse will tell you when to have these blood tests. Tell your doctor if you are on an anticoagulant (medication used to treat or prevent blood clots) e.g. warfarin. You may need to have additional blood tests.

### Pumps and central venous access devices (CVADs)

This treatment involves having chemotherapy through a pump. To have this, you will also need a central venous access device (CVAD). Your doctor or nurse will explain this to you. For more information see the [eviQ patient information sheets on pumps and CVADs](#). At home you will need to look at your pump 3 to 4 times a day to check it is working. Your nurse will teach you how to do this.

### Other medications given during this treatment

- **Anti-sickness (anti-nausea) medication:** you may be given some anti-sickness medication. Make sure you take this medication as your doctor or nurse tells you, even if you don't feel sick. This can help to prevent the sickness starting.
- **Antidiarrhoeals:** you will be given some medication called loperamide to treat the diarrhoea. See the *Side effects* section below for further information about diarrhoea and for instructions on how and when to take the loperamide.
- **Medication for skin rash:** you may be given some medication (which may include a steroid cream, an antibiotic cream and/or oral antibiotics) to prevent or treat skin rash. Your doctor or nurse will tell you how to take or use these medications.
- **Cetuximab premedication:** before your treatment with cetuximab you will need to take some tablets called a premedication to help prevent you from having a reaction to the cetuximab.

## Side effects

Cancer treatments can cause damage to normal cells in your body, which can cause side effects. Everyone gets different side effects, and some people will have more problems than others.

The table below shows some of the side effects you may get with this treatment. You are unlikely to get all of those listed and you may also get some side effects that have not been listed.

Tell your doctor or nurse about any side effects that worry you. Follow the instructions below and those given to you by your doctor or nurse.

Immediate (onset hours to days)	
<b>Allergic reaction</b>	<ul style="list-style-type: none"> <li>• Allergic reactions are uncommon but can be life threatening.</li> <li>• <b>If you feel unwell during the infusion or shortly after it, or:</b> <ul style="list-style-type: none"> <li>◦ <b>get a fever, shivers or shakes</b></li> <li>◦ <b>feel dizzy, faint, confused or anxious</b></li> <li>◦ <b>start wheezing or have difficulty breathing</b></li> <li>◦ <b>have a rash, itch or redness of the face</b></li> </ul> </li> </ul> <p><b><u>While you are in hospital:</u> Tell your doctor or nurse immediately.</b></p> <p><b><u>After you leave:</u> Contact your doctor or nurse immediately, or go to the nearest hospital Emergency Department.</b></p>
<b>Diarrhoea (early onset)</b>	<ul style="list-style-type: none"> <li>• You may get bowel motions (stools, poo) that are more frequent or more liquid (diarrhoea).</li> <li>• You may also get: <ul style="list-style-type: none"> <li>◦ bloating, cramping or pain</li> <li>◦ increased saliva, a runny nose or watery eyes</li> <li>◦ sweating or flushing.</li> </ul> </li> <li>• These symptoms are caused by the drug irinotecan. They can occur during or shortly after the drug has been given.</li> <li>• <b>Tell your doctor or nurse immediately if you develop any of these symptoms.</b></li> </ul>
<b>Nausea and vomiting</b>	<ul style="list-style-type: none"> <li>• You may feel sick (nausea) or be sick (vomit).</li> <li>• Take your anti-sickness medication as directed even if you don't feel sick.</li> <li>• Drink plenty of fluids (unless you are fluid restricted).</li> <li>• Eat small meals more frequently.</li> <li>• Try food that does not require much preparation.</li> <li>• Try bland foods like dry biscuits or toast.</li> <li>• Gentle exercise may help with nausea.</li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Nausea and vomiting during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have uncontrolled vomiting or feel dizzy or light-headed.</b></li> </ul>
<b>Taste and smell changes</b>	<ul style="list-style-type: none"> <li>• You may find that food loses its taste or tastes different.</li> <li>• These changes are likely to go away with time.</li> <li>• Do your mouth care regularly.</li> <li>• Chew on sugar-free gum or eat sugar-free mints.</li> <li>• Add flavour to your food with sauces and herbs.</li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Taste and smell changes during cancer treatment</a>.</li> </ul>
<b>Heart problems</b>	<ul style="list-style-type: none"> <li>• You may get: <ul style="list-style-type: none"> <li>◦ chest pain or tightness</li> <li>◦ shortness of breath</li> <li>◦ an abnormal heartbeat</li> </ul> </li> <li>• Tell your doctor if you have a history of heart problems or high blood pressure.</li> <li>• <b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the symptoms listed above.</b></li> </ul>

**Early (onset days to weeks)**

<b>Infection risk (neutropenia)</b>	<ul style="list-style-type: none"> <li>• This treatment lowers the amount of white blood cells in your body. The type of white blood cells that help to fight infection are called neutrophils. Having low level of neutrophils is called neutropenia. If you have neutropenia, you are at greater risk of getting an infection. It also means that your body can't fight infections as well as usual. This is a serious side effect, and can be life threatening.</li> <li>• Wash your hands often.</li> <li>• Keep a thermometer at home and take your temperature regularly, and if you feel unwell.</li> <li>• Do your mouth care regularly.</li> <li>• Inspect your central line site (if you have one) daily for any redness, pus or swelling.</li> <li>• Limit contact with people who are sick.</li> <li>• Learn how to recognise the signs of infection.</li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Infection during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the following signs or symptoms:</b> <ul style="list-style-type: none"> <li>◦ <b>a temperature of 38°C or higher</b></li> <li>◦ <b>chills, shivers, sweats or shakes</b></li> <li>◦ <b>a sore throat or cough</b></li> <li>◦ <b>uncontrolled diarrhoea</b></li> <li>◦ <b>shortness of breath</b></li> <li>◦ <b>a fast heartbeat</b></li> <li>◦ <b>become unwell even without a temperature.</b></li> </ul> </li> </ul>
<b>Low platelets (thrombocytopenia)</b>	<ul style="list-style-type: none"> <li>• This treatment lowers the amount of platelets in your blood. Platelets help your blood to clot. When they are low, you are at an increased risk of bleeding and bruising.</li> <li>• Try not to bruise or cut yourself.</li> <li>• Avoid contact sport or vigorous exercise.</li> <li>• Clear your nose by blowing gently.</li> <li>• Avoid constipation.</li> <li>• Brush your teeth with a soft toothbrush.</li> <li>• Don't take aspirin, ibuprofen or other similar anti-inflammatory medications unless your doctor tells you to.</li> <li>• Tell your doctor or nurse if you have any bruising or bleeding.</li> <li>• <b>Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if you have any uncontrolled bleeding.</b></li> </ul>
<b>Diarrhoea (late onset)</b>	<ul style="list-style-type: none"> <li>• You may get bowel motions (stools, poo) that are more frequent or more liquid (diarrhoea).</li> <li>• You may also get bloating, cramping or pain.</li> <li>• These symptoms are caused by the drug irinotecan. This can become very serious and life threatening if not treated quickly and properly.</li> <li>• Take your antidiarrhoea medication, loperamide, as prescribed: When the diarrhoea starts, take 4 mg (this is 2 tablets/capsules), then take one tablet/capsule (2 mg) every 2 hours during the day and 2 tablets/capsules (4 mg) every 4 hours at night while you still have diarrhoea and until the diarrhoea has stopped for 12 hours. You should not take loperamide at these doses for more than 48 hours.</li> <li>• Drink plenty of fluids (unless you are fluid restricted).</li> <li>• Eat and drink small amounts more often.</li> <li>• Avoid spicy foods, dairy products, high fibre foods, and coffee.</li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Diarrhoea during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> <li>• <b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have diarrhoea for more than 24 hours, any blood in your bowel motions, or a temperature of 38°C or higher.</b></li> </ul>
<b>Appetite loss (anorexia)</b>	<ul style="list-style-type: none"> <li>• You may not feel like eating.</li> <li>• Try to avoid drinking fluids at meal times.</li> <li>• Try to eat small meals or snacks regularly throughout the day.</li> <li>• Try to eat food that is high in protein and calories.</li> <li>• If you are worried about how much food you can eat, or if you are losing weight, ask to speak to a dietitian.</li> </ul>

<b>Mouth pain and soreness (mucositis)</b>	<ul style="list-style-type: none"> <li>• You may have: <ul style="list-style-type: none"> <li>◦ bleeding gums</li> <li>◦ mouth ulcers</li> <li>◦ a white coating on your tongue</li> <li>◦ pain in the mouth or throat</li> <li>◦ difficulty eating or swallowing.</li> </ul> </li> <li>• Avoid spicy, acidic or crunchy foods and very hot or cold food and drinks.</li> <li>• Try bland and soft foods.</li> <li>• Brush your teeth gently with a soft toothbrush after each meal and at bedtime. If you normally floss continue to do so.</li> <li>• Rinse your mouth after you eat and brush your teeth, using either: <ul style="list-style-type: none"> <li>◦ 1/4 teaspoon of salt in 1 cup of warm water, or</li> <li>◦ 1/4 teaspoon of bicarbonate of soda in 1 cup of warm water</li> </ul> </li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Mouth problems during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>
<b>Tiredness and lack of energy (fatigue)</b>	<ul style="list-style-type: none"> <li>• You may feel very tired, have no energy, sleep a lot, and not be able to do normal activities or things you enjoy.</li> <li>• Do not drive or operate machinery if you are feeling tired.</li> <li>• Nap for short periods (only 1 hour at a time)</li> <li>• Prioritise your tasks to ensure the best use of your energy.</li> <li>• Eat a well balanced diet and drink plenty of fluids (unless you are fluid restricted).</li> <li>• Try some gentle exercise daily.</li> <li>• Allow your friends and family to help.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>
<b>Eye problems</b>	<ul style="list-style-type: none"> <li>• You may get: <ul style="list-style-type: none"> <li>◦ eye pain</li> <li>◦ red, sore or swollen eyes</li> <li>◦ blurred vision</li> <li>◦ watery or gritty eyes</li> <li>◦ changes in your eyesight</li> <li>◦ sensitivity to sunlight.</li> </ul> </li> <li>• Protect your eyes from the weather (sun and wind) by wearing sunglasses, especially if you have lost your eyelashes.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above. Eye drops may help with your symptoms.</b></li> </ul>
<b>Skin rash (acneiform rash)</b>	<ul style="list-style-type: none"> <li>• You may get an acne-like skin rash.</li> <li>• Your skin may become red and dry.</li> <li>• Moisturise your skin with a gentle non-perfumed moisturising cream like sorbolene or aqueous cream.</li> <li>• Do not scratch your skin.</li> <li>• Do not use over-the-counter acne treatments as these can make the rash worse.</li> <li>• Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and sunscreen of SPF 50 or higher.</li> <li>• You may be given medications to prevent the rash.</li> <li>• <b>Tell your doctor or nurse as soon as possible if you notice any changes to the rash like itching, pain or pus forming</b></li> </ul>
<b>Skin changes</b>	<ul style="list-style-type: none"> <li>• Your skin may become dry, and you may notice changes to areas of your skin that have been exposed to the sun.</li> <li>• Keep your skin moisturised with a cream such as sorbolene or aqueous cream.</li> <li>• Avoid direct sunlight.</li> <li>• Protect your skin from the sun by wearing a wide-brimmed hat, sun-protective clothing, sunglasses and sunscreen of SPF 50 or higher.</li> <li>• <b>Tell your doctor or nurse if you notice any skin changes.</b></li> </ul>

<b>Skin that is more sensitive to the sun (photosensitivity)</b>	<ul style="list-style-type: none"> <li>• After being out in the sun you may develop a rash like a bad sunburn.</li> <li>• Your skin may become red, swollen and blistered.</li> <li>• Avoid direct sunlight.</li> <li>• Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and a sunscreen of SPF 50 or higher.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>
<b>Hand-foot syndrome (palmar-plantar erythrodysesthesia)</b>	<ul style="list-style-type: none"> <li>• The palms of your hands and soles of your feet may become: <ul style="list-style-type: none"> <li>◦ red and hot</li> <li>◦ swollen</li> <li>◦ painful and tender</li> <li>◦ blistered.</li> </ul> </li> <li>• The skin in the area may also peel.</li> <li>• Moisturise your hands and feet daily with sorbolene or aqueous cream.</li> <li>• Keep your hands and feet clean and dry.</li> <li>• Avoid hot water, instead use lukewarm water to bathe.</li> <li>• Avoid direct sunlight.</li> <li>• Avoid unnecessary walking, jogging or exercise.</li> <li>• Wear cotton socks and avoid tight-fitting shoes.</li> <li>• <b>Tell your doctor or nurse as soon as possible if you notice any skin changes on your hands or feet.</b></li> </ul>

Late (onset weeks to months)	
<b>Low red blood cells (anaemia)</b>	<ul style="list-style-type: none"> <li>You may feel dizzy, light-headed, tired and appear more pale than usual.</li> <li>Tell your doctor or nurse if you have any of these signs or symptoms. You might need a blood transfusion.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have any chest pain, trouble breathing, or feel like your heart is racing.</b></li> </ul>
<b>Swelling and pain around the fingernails or toenails (paronychia)</b>	<ul style="list-style-type: none"> <li>The skin around your nails may swell and become painful.</li> <li>Apply a warm compress or soak your nails for 15 minutes, 3 or 4 times a day, in warm water or a mixture of equal parts vinegar and water.</li> <li>Keep your nails clean and short.</li> <li>Avoid things like biting your fingernails, getting a manicure, pedicure or false nails.</li> <li>Wear gloves when you wash the dishes, work in the garden, or clean the house.</li> <li>Tell your doctor or nurse if you get any of the symptoms listed above.</li> </ul>
<b>Hair changes</b>	<ul style="list-style-type: none"> <li>Your hair may become fine or curly and may break easily.</li> <li>Your eyelashes and eyebrows may grow more than normal.</li> <li>Use a gentle shampoo and a soft hairbrush.</li> <li>Take care with hair products like hairspray, hair dye, bleaches and perms.</li> <li>Ask your doctor or nurse about the <a href="http://www.lgfb.org.au">Look Good Feel Better</a> program (www.lgfb.org.au).</li> </ul>
<b>Skin colour changes</b>	<ul style="list-style-type: none"> <li>You may have darkening of your skin, especially in areas that are exposed to the sun.</li> <li>You may also notice darkening of your tongue, gums and over your finger joints.</li> <li>These skin changes may fade over time.</li> <li>Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and a sunscreen of SPF 50 or higher.</li> </ul>
<b>Lung problems</b>	<ul style="list-style-type: none"> <li>Lung problems are rare, but can be serious. They may occur throughout treatment or after the completion of treatment.</li> <li>You may get: <ul style="list-style-type: none"> <li>shortness of breath</li> <li>fever</li> <li>dry cough</li> <li>wheezing</li> <li>fast heartbeat</li> <li>chest pain.</li> </ul> </li> <li>Your doctor will monitor how well your lungs are working during your treatment.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have chest pain or become short of breath.</b></li> </ul>

## General advice for people having cancer treatment

### Chemotherapy safety

- Learn how to keep you and your family safe while you are having anticancer drugs.
- See our patient information sheet - [Chemotherapy safety at home](#).

### Blood clot risk

- Cancer and anticancer drugs can increase the risk of a blood clot (thrombosis).
- Tell your doctor if you have a family history of blood clots.
- A blood clot can cause pain, redness, swelling in your arms or legs, shortness of breath or chest pain.
- If you have any of these symptoms go to your nearest hospital Emergency Department.

### Medications and vaccinations

- Before you start treatment, tell your doctor about any medications you are taking, including vitamins or herbal supplements.
- Don't stop or start any medications during treatment without talking to your doctor and pharmacist first.



- Paracetamol is safe to take if you have a headache or other mild aches and pains. It is recommended that you avoid taking aspirin, ibuprofen and other anti-inflammatory type medications for pain while you are having treatment. However, if these medications have been prescribed by your doctor, do not stop taking them without speaking with your doctor.
- Vaccinations such as flu and tetanus vaccines are safe to receive while having treatment. Do not have any live vaccines during your treatment or for 6 months after it finishes. If you are unsure, check with your doctor before you have any vaccinations.
- People you live with should be fully vaccinated, including having live vaccines according to the current vaccination schedule. Extra care needs to be taken with hand washing and careful disposal of soiled nappies for infants who have recently received the rotavirus vaccine.

### Other medical and dental treatment

- If you go to hospital or any other medical appointment (including dental appointments), always tell the person treating you that you are receiving anticancer drugs.
- Before you have any dental treatment, talk to your doctor.

### Diet

- While you are receiving this treatment it is important that you try to maintain a healthy diet.
- Grapefruit and grapefruit juice can interact with your medication and should be avoided while you are on this treatment.
- Speak to your doctor or nurse about whether drinking alcohol is safe with your treatment.
- If you have any concerns about recent weight loss or weight gain or questions about your diet, ask to speak to a dietitian.

### Fertility

- Some cancer treatments can reduce your fertility. This can make it difficult or impossible to get pregnant or father a child.
- Talk to your doctor or nurse before you start any treatment. Depending on your situation there may be fertility sparing options available to you and/or your partner, discuss these with your doctor or nurse.

### Pregnancy and breastfeeding

- Some cancer treatments can be dangerous to unborn babies. Talk to your doctor or nurse if you think there is any chance that you could be pregnant.
- Do not try to get pregnant or father a child during this treatment. Contraception should be used during treatment and after stopping treatment. Ask your doctor or nurse about what type of contraception you should use.
- If you are planning pregnancy/fatherhood after completing this treatment, talk to your doctor. Some doctors advise waiting between 6 months and 2 years after treatment.
- Do not breastfeed if you are on this treatment, as anti-cancer medications can also pass into breast milk.

### Sex life and sexuality

- The desire to have sex may decrease as a result of this treatment or its side effects.
- Your emotions and the way you feel about yourself may also be affected by this treatment.
- It may help to discuss your concerns with your partner and doctor or nurse.

### Quitting smoking

- It is never too late to quit smoking. Quitting smoking is one of the best things you can do to help your treatment work better.
- There are many effective tools to improve your chances of quitting.
- Talk to your treating team for more information and referral to a smoking cessation support service.

### Staying active

- Research shows that exercise, no matter how small, has many benefits for people during and after cancer treatment.
- Talk to your doctor before starting an exercise program. Your doctor can advise whether you need a modified exercise program.

**For more information about cancer treatment, side effects and side effect management see our [Patient and carers](#) section.**

## Where to get more information

### Telephone support

- Call Cancer Council on 13 11 20 for cancer information and support

### Bowel cancer information

- Australian Council of Stoma Associations – [australianstoma.com.au](http://australianstoma.com.au)
- Australian Government Bladder and Bowel – [bladderbowel.gov.au](http://bladderbowel.gov.au)
- Australian Government Department of Health & Ageing Stoma appliance scheme – [health.gov.au/internet/main/publishing.nsf/Content/Stoma+Appliance+Scheme-1](http://health.gov.au/internet/main/publishing.nsf/Content/Stoma+Appliance+Scheme-1)
- Bowel Cancer Australia – [bowelcanceraustralia.org](http://bowelcanceraustralia.org)
- National Public Toilet map – [toiletmap.gov.au](http://toiletmap.gov.au)
- Recovering after Pelvic Radiation Therapy: A guide for women – <https://www.targetingcancer.com.au/useful-resources/recovering-after-pelvic-radiation-therapy-a-guide-for-women/>

**General cancer information and support**

- Australian Rare Cancer (ARC) Portal – [arcportal.org.au/](http://arcportal.org.au/)
- Beyondblue – [beyondblue.org.au](http://beyondblue.org.au)
- Cancer Australia – [canceraustralia.gov.au](http://canceraustralia.gov.au)
- Cancer Council Australia – [cancer.org.au](http://cancer.org.au)
- Cancer Voices Australia – [cancervoicesaustralia.org](http://cancervoicesaustralia.org)
- CanTeen – [canteen.org.au](http://canteen.org.au)
- Carers Australia – [carersaustralia.com.au](http://carersaustralia.com.au)
- CHILL Cancer related hair loss - [scalpcooling.org](http://scalpcooling.org)
- eviQ Cancer Treatments Online – [eviQ.org.au](http://eviQ.org.au)
- LGBTQI+ People and Cancer - [cancerCouncil.com.au/cancer-information/lgbtqi](http://cancerCouncil.com.au/cancer-information/lgbtqi)
- Look Good Feel Better – [lgfb.org.au](http://lgfb.org.au)
- Patient Information – [patients.cancer.nsw.gov.au](http://patients.cancer.nsw.gov.au)
- Radiation Oncology Targeting Cancer – [targetingcancer.com.au](http://targetingcancer.com.au)
- Redkite – [redkite.org.au](http://redkite.org.au)
- Return Unwanted Medicines – [returnmed.com.au](http://returnmed.com.au)
- Staying active during cancer treatment – [patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active](http://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active)

**Quit smoking information and support**

Quitting smoking is helpful even after you have been diagnosed with cancer. The following resources provide useful information and support to help you quit smoking. Talk to your treating team about any other questions you may have.

- Call Quitline on 13 QUIT (13 78 48)
- iCanQuit – [iCanQuit.com.au](http://iCanQuit.com.au)
- Patient Information - [patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking](http://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking)
- Quitnow – [quitnow.gov.au](http://quitnow.gov.au)

**Additional notes:**

---

---

---

---

---

---

---

---

---

---

---

This document is a guide only and cannot cover every possible situation. The health professionals caring for you should always consider your individual situation when making decisions about your care. Contact your cancer clinic staff or doctor if you have any questions or concerns about your treatment, or you are having problems coping with side effects. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use of this document is subject to eviQ's disclaimer available at [www.eviQ.org.au](http://www.eviQ.org.au)

---

**First approved:** 27 April 2015

**Last reviewed:** 20 October 2022

**Review due:** 31 December 2024

***The currency of this information is guaranteed only up until the date of printing, for any updates please check:***

<https://www.eviq.org.au/pi/1682>

12 Jul 2023