

P  
R  
E  
V  
E  
N  
T  
I  
O  
N

**AIM:** clean, dry, intact, comfortable, cost-effective dressing that provides catheter or TIVAD needle securement.

**KEY PRACTICES**

- **Assess and document dressing:** at least once per shift, and each outpatient, clinic or home visit
- **IV administration lines:** ensure IV lines are secured to patient to prevent pull on the dressing
- **Replace dressing every 7 days:** or earlier if it contains ooze, moisture or is not intact

**PATIENT FACTORS**

- **Educate patient** to look at dressing each day, to notify nurse of discomfort, burning, or itch of skin under dressing
- **Encourage hydration and balanced diet:** important elements for skin health
- **Hair under dressing:** clip if required to assist with dressing adherence to skin



Photo permission: E Alexandrou

**Types of CVADs**

- **CICC:** centrally inserted central catheter
- **tc-CICC:** tunnelled cuffed CICC (central venous catheter)
- **PICC:** peripherally inserted central catheter
- **TIVAD:** totally implantable venous access device (portacath)
- **Apheresis catheters**

Photo permission: Interrad Medical

S  
T  
A  
N  
D  
A  
R  
D  
C  
A  
R  
E

**KEY PRACTICE POINTS**

**SKIN ANTISEPSIS**

- ✓ **2% CHG in 70% IPA swab sticks**
  - e.g. Chloraprep® (sterile applicator), 3M™ Soluprep™ SwabStick, Reynard Foam Swab Stick®
- ✓ **Prolonged skin antiseptics**
  - CHG disc e.g. Biopatch® OR
  - CHG gel dressing 3M™ Tegaderm™ CHG Dressing
- ✓ **Tissue adhesive at catheter exit site**
  - (bacteriostatic, haemostatic) e.g. SecurePortIV®
- ✓ **Alternatives: silver or PMHB discs**
  - e.g. Acticoat™ Site, Kendall™ AMD Antimicrobial Foam Discs
- ! **CRITICAL PRACTICE POINT**  
Allow skin to completely air dry before applying any dressing materials

**DRESSING MATERIAL**

- ✓ **Application:** do not stretch or apply with tension, ensure borders adhere to the skin
- ✓ **Use sterile TSM:** consider next generation fabric bordered dressings
  - **Bordered TSM with integrated securement** e.g. SorbaView Shield® Dressing
  - **Bordered TSM** e.g. 3M™ Tegaderm™ I.V. Advanced Securement Dressing
- ! **CRITICAL PRACTICE POINT**  
Use low and slow, controlled removal technique while supporting skin to prevent skin damage related to acrylic adhesives\*
- \* **Acrylic adhesive:** strong dressing adhesive that increases risk of skin damage during removal

**SKIN PROTECTION**

- ✓ **Use alcohol-free skin barrier film** under dressing materials (not under CHG gel or disc) e.g. Cavilon™ No Sting Barrier Film
  - Preventatively for any patients with **risk factors for skin impairment** \*\*
  - For diaphoretic patients for dressing adhesion
- ! **CRITICAL PRACTICE POINT**  
Allow skin to completely air dry before applying any dressing materials

**SECUREMENT**

- ✓ **Use engineered securement devices (ESD)**
  - **Adhesive ESD** e.g. StatLock®, GripLoc®, bordered TSM with silicone adhesive ESD e.g. 3M™ PICC-CVC CHG Securement Dressing
  - **Subcutaneous ESD:** no adhesives for use with impaired skin e.g. SecurAcath™
- ✓ **Tissue adhesive** applied to catheter exit site (haemostatic, bacteriostatic, securement) and potentially under catheter e.g. SecurePortIV®
- ! **CRITICAL PRACTICE POINT**  
All external catheters are secured at all times ideally within 1-2cm of exit site

**\*\* RISK FACTORS FOR SKIN IMPAIRMENT**

- **Age** >50 years, neonates, premature infants
- **Number & type of comorbidities:** infection, diabetes, renal failure, malignancy, skin conditions e.g. eczema
- **Medication:** anticancer agents e.g. metabolites (e.g. fluorouracil, capecitabine), taxanes (e.g. paclitaxel); angiogenesis inhibitors (e.g. bevacizumab); epidermal growth factor inhibitors (e.g. cetuximab); long-term corticosteroids
- **Radiation therapy**
- **Immunosuppression**
- **Malnutrition & dehydration**

Continue with standard care

NO

SKIN CHANGES

YES

**ABBREVIATIONS**

- CHG - chlorhexidine
- ESD - engineered securement device (for catheter securement)
- IPA - isopropyl alcohol
- IV - intravenous
- NaCl - sodium chloride
- NC - needleless connector
- PHMB - polyhexamethylene biguanide
- TSM - transparent semipermeable membrane

**COMPLETE HOLISTIC PATIENT ASSESSMENT INCLUDING:**

- Patient diagnosis
- Comorbidities
- Prescribed and other treatments
- Allergies and sensitivities
- Existing skin conditions
- Cognitive performance

! Consult vascular access or wound care specialist nurse if required

Skin changes can be categorised as infection, dermatitis or mechanical injury related

A  
S  
S  
E  
S  
S  
M  
E  
N  
T

**LOCAL SKIN INFECTION**

**Signs / Symptoms**

- Red, blanched, mottled, warmth, discomfort, oedema, pustules, exudate, odour, induration

**Assessment**

- Is the patient neutropenic?
- Is the dressing clean, dry and intact?
- Is the skin broken? Evidence of skin impairment?
- What is the amount and type of exudate present?

**Key Points**

- ✓ Discuss assessment with treating medical team
- ✓ As per orders - skin culture of exit site, systemic antibiotics, potential CVAD removal
- ✓ Continue regular observation, patient assessment and documentation




Photo permissions: K Curtis

**CONTACT DERMATITIS**

- irritant, allergic, moisture-associated

**Signs / Symptoms**

- Redness, burning, pain, itch, shiny, scales, vesicles, exudate (blood, serous), maceration (moisture-associated)

**Assessment**

- Is it associated with infiltration or extravasation?
- Is the area the same size & shape of dressing material or adhesive ESD?
- What are suspected cause/s - solutions, dressing materials, adhesives?
- What is the amount and type of exudate present?
- Is the skin broken?

**Key Points**

- Discuss assessment with the vascular access or wound care specialist nurse if required



Photo permissions: K Curtis

**MECHANICAL INJURY**

**Signs / Symptoms**

- Initial skin injury from insertion (bleeding exit site), skin stripping, tears, tension blisters, pressure injury, bruising

**Assessment**

- What caused the injury?
- Skin tear: can the skin edges be gently moved together?
- Was the pressure injury caused by a bandage, tubing, needleless connector?


**Key Points**

- Avoid the mechanism of injury
- Protect skin from catheter hubs or NC with cushioning dressing materials, avoid tight tubing material over dressing



Photo permissions: K Curtis

! **PROMPT INTERVENTION:** Do not wait until the next dressing change is due



## SKIN ANTISEPSIS ALTERNATIVES

- ✓ Remove exudate with sterile NaCl
- ✓ Continue with 2% CHG in 70% IPA swab sticks
- ✓ Ensure correct application of CHG disc (blue side up, 360° coverage around catheter at the exit site), gel pad covering exit site

**CRITICAL PRACTICE POINT**  
Allow skin to completely air dry before applying any dressing materials

- Identify and replace suspected irritant/s - may be a combination including:
- ✓ CHG: skin patch test\*\*\* with reduced CHG concentration (0.5%) or
  - ✓ Alcohol: change to aqueous CHG
  - ✓ CHG sensitivity: skin patch test\*\*\* with povidone-iodine 10%
  - ✓ Broken skin: use aqueous CHG (avoid pain caused by alcohol on broken skin)
  - ✓ Use sterile NaCl: for severe skin impairment - allow for greater drying time
- \*\*\* Skin patch test  
Apply material or solution to skin (e.g. forearm)  
Monitor for short period (30-60 mins)  
Monitor over next 2-3 days  
Observe for signs, ask patient re symptoms of irritation

- ✓ Continue to use 2% CHG in 70% IPA swab sticks if skin is intact
- ✓ Consider temporary reduction to CHG 0.5% in 70% IPA or aqueous until skin is healed
- ✓ If severe use sterile NaCl for maintenance procedures - allow for greater drying time

**CRITICAL PRACTICE POINT**  
Allow greater drying time for NaCl & povidone-iodine solutions

## DRESSING MATERIAL ALTERNATIVES

- ✓ CHG gel bordered TSM: allows observation of exit site
- ✓ Antimicrobial disc: add to dressing management if not currently used e.g. CHG (Biopatch® or gel dressing 3M™ Tegaderm™ CHG Dressing), silver (Acticoat™ Site) or PHMB discs (Kendall™ AMD Antimicrobial Foam Discs)
- ✓ For exudate: use absorbent dressing material and attempt to keep separate from exit site e.g. Mepilex® Border, Mepilex Lite®, Biatain® Non-adhesive dressings

- ✓ Identify and replace suspected irritant/s: may be more than one agent
- ✓ Use NaCl or alcohol-free adhesive removal products (e.g. wipes, sprays, pads) to remove dressing. Cleanse skin with sterile NaCl to remove product from skin before applying skin antiseptis
- ✓ For exudate: use absorbent dressing material and attempt to keep separate from exit site e.g. Mepilex® Border, Mepilex Lite®, Biatain® Non-adhesive dressings
- ✓ Consider changing TSM brand e.g. IV3000™, Opsite Flexigrid™, SorbaView Shield® Dressing, 3M™ Tegaderm™ I.V. Advanced Securement Dressing
- ✓ Ensure impaired skin is protected and completely covered
- ✓ Slightly move dressing materials to avoid applying adhesive to same skin area every dressing

- ✓ Continue to use 2% CHG in 70% IPA swab sticks if skin is intact
- ✓ Consider temporary reduction to CHG 0.5% in 70% IPA or aqueous until skin is healed
- ✓ If severe use sterile NaCl for maintenance procedures - allow for greater drying time

**CRITICAL PRACTICE POINT**  
Minimise skin damage with dressing removal technique. Use low and slow technique (for bordered dressings), or lateral stretch technique for flat dressings while supporting the skin (IV3000™)

## SKIN PROTECTION ALTERNATIVES

- ✓ Avoid use of hydrocolloid materials\*\*\*\*
- ✓ Unbroken skin: use alcohol-free skin barrier film e.g. Cavilon™ No Sting Barrier Film
- ✓ Broken skin: use alcohol-free skin barrier film designed for broken skin e.g. 3M™ Cavilon™ Advanced Skin Protectant
- ✓ Use silicone adhesive materials\*\*\*\*\* for protection: over impaired skin to protect from TSM, adhesive ESD

**Main types of adhesives used for dressing materials:**

- \* Acrylic adhesive: strong adhesive for dressing adherence; greater risk of skin damage during removal so slow and low, controlled removal technique is required possibly with additional adhesive remover wipes
- \*\*\*\* Hydrocolloid: avoid due to moisture content under dressing
- \*\*\*\*\* Silicone adhesive: gentle adhesive; can be applied over impaired skin; some materials have absorbent properties to wick away excess moisture; gentle on impaired skin for removal

- ✓ If skin is intact: continue to use skin barrier film
- ✓ If skin is broken: apply
  - silicone adhesive material as protective layer between skin and TSM adhesive e.g. Mepilex® Border, Mepilex Lite®, Biatain® Non-adhesive dressing
  - skin barrier film designed for broken skin e.g. 3M™ Cavilon™ Advanced Skin Protectant
- ✓ Severe cases: place silicone adhesive dressing material over broken skin and situate all dressing materials on top (see example below)
- ✓ Skin with high moisture content or ooze: consider using silicone adhesive material with absorbent properties e.g. Mepilex Border™ or Biatain® Silicone dressing

- ✓ If skin intact: continue to use skin barrier film and ensure skin is completely air dry before dressing application
- ✓ If skin is broken: apply silicone material (protection layer between skin and TSM adhesive)
- ✓ Severe cases: use silicone adhesive dressing material on the skin and place all dressing materials on top
- ✓ Skin with high moisture content or ooze: consider using silicone adhesive material with absorbent properties
- ✓ Broken skin: cover with silicone dressing material

**CRITICAL PRACTICE POINT**  
Allow skin to completely air dry before applying any dressing materials

## SECUREMENT ALTERNATIVES

- ✓ Adhesive ESD: ensure it is not over or adjacent to catheter exit site for monitoring
- ✓ Use a subcutaneous ESD: advantage is there is no adhesive involved, can place silicone adhesive material underneath if needed

- ✓ Move adhesive ESD to different area of skin to avoid skin impairment
- ✓ Place silicone adhesive material between skin and adhesive or subcutaneous ESD
- ✓ Use subcutaneous ESD

- ✓ Alternative: bordered dressing that includes ESD with silicone adhesive
- ✓ Move adhesive ESD to avoid injured skin
- ✓ Use a subcutaneous ESD, can place silicone adhesive dressing underneath if required

CONTINUE TO ASSESSEVERY DRESSING. DOCUMENTALL ASSESSMENTS, PROCEDURES & INTERVENTIONS

### EXAMPLE



Figure 1: Erosive contact dermatitis




Figure 2: Impaired skin protected with silicone adhesive dressing with adhesive ESD on top, and completely covered with TSM



Figure 3: One week later



**References:**

1. Australian and New Zealand College of Anaesthetists. (2018). *Topical application of chlorhexidine and the risks of accidental injection in regional anaesthesia and vascular access procedures*. Retrieved from <http://www.anzca.edu.au/documents/8-2-2-anzca-and-acsghc-joint-statement-on-chlorhex.pdf>:
2. Broadhurst, D., Moureau, N. L., & Ullman, A. J. (2017). Management of Central Venous Access Device-Associated Skin Impairment. An Evidence-Based Algorithm. *J Wound Ostomy Continence Nurs*, *44*(3), 1-10.
3. Cartagena, G. (2018). *Skin Matters: Impact of Hospital-Acquired Skin Injuries in the Adult Patient*. Paper presented at the World Congress on Vascular Access, Denmark, Copenhagen.
4. Chan RJ, Northfield S, Larsen E, Mihala G, Ullman A, Hancock P, et al. Central venous Access device SeCurement And Dressing Effectiveness for peripherally inserted central catheters in adult acute hospital patients (CASCADE): a pilot randomised controlled trial. *Trials*. 2017;18(1):458.
5. Corley, A., Marsh, N., Ullman, A. J., & Rickard, C. M. (2017). Tissue adhesive for vascular access devices: who, what, where and when? *Br J Nurs*, *26*(19), S4-s17. doi:10.12968/bjon.2017.26.19.S4
6. Hitchcock, J., & Savine, L. (2017). Medical adhesive-related skin injuries associated with vascular access. *Br J Nurs*, *26*(8), S4-s12. doi:10.12968/bjon.2017.26.8.S4
7. Kutzscher, L. (2012). Management of Irritant Contact Dermatitis and Peripherally Inserted Central Catheters. *Clinical Journal of Oncology Nursing*, *16*(2), E48-E55. doi:10.1188/12.CJON.E48-E55
8. McNichol, L., Lund, C., Rosen, T., & Gray, M. (2013). Medical adhesives and patient safety: state of the science: consensus statements for the assessment, prevention, and treatment of adhesive-related skin injuries. *Journal of Wound, Ostomy & Continence Nursing*, *40*(4), 365-380. doi:10.1097/won.0b013e3182995516
9. Mitchell ML, Ullman AJ, Takashima M, Davis C, Mihala G, Powell M, et al. Central venous access device Securement and dressing effectiveness: The CASCADE pilot randomised controlled trial in the adult intensive care. *Aust Crit Care*. 2020;33(5):441-51.
10. Ralph Webber, J. L., & Maningo-Salinas, M. J. (2020). "Sticking It to Them"—Reducing Migration of Peripherally Inserted Central Catheters. *Journal of the Association for Vascular Access*, *25*(1), 10-15. doi:10.2309/j.java.2020.001.004
11. Rickard, C., Ullman, A., Kleidon, T., & Marsh, N. (2017). TEN TIPS FOR DRESSING AND SECUREMENT OF IV DEVICE WOUNDS. *Australian Nursing & Midwifery Journal*, *24*(10), 32-34.
12. Ullman, A. J., Cooke, M., & Rickard, C. M. (2015). Examining the Role of Securement and Dressing Products to Prevent Central Venous Access Device Failure: A Narrative Review. *Journal of the Association for Vascular Access*, *20*(2), 99-110. doi:10.1016/j.java.2015.03.001
13. Webster, J., Larsen, E., Marsh, N., Choudhury, A., Harris, P., & Rickard, C. M. (2017). Chlorhexidine gluconate or polyhexamethylene biguanide disc dressing to reduce the incidence of central-line-associated bloodstream infection: a feasibility randomized controlled trial (the CLABSI trial). *Journal of Hospital Infection*, *96*(3), 223-228. doi:<https://dx.doi.org/10.1016/j.jhin.2017.04.009>