

Testicular germ cell seminoma adjuvant cARBOplatin

ID: 325 v.4 **Endorsed** Essential Medicine List

⚠️ ADDIKD Carboplatin dosing:

For dosing carboplatin, ADDIKD recommends that:

- Directly measured glomerular filtration rate (mGFR) is the preferred kidney function value in the Calvert formula, especially where estimated kidney function may be unreliable for accurate therapeutic dosing.
- Where mGFR is unavailable, eGFR adjusted to an individual's body surface area (BSA-adjusted eGFR) is a suitable alternative for use in the Calvert formula.
- Kidney function should not be capped at 125 mL/min for use in the Calvert formula.
- Recalculation of carboplatin doses at each cycle is unnecessary, except when baseline kidney function (e.g., eGFR) alters by > 20% or when there is a change in the clinical status of the patient.

For further information refer the [eviQ Factsheet](#) around carboplatin dosing and the carboplatin drug monograph within the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\)](#) and [carboplatin dose calculators](#).

[Link to ANZUP testicular cancer surveillance recommendations](#)

Check for clinical trials in this patient group. [Link to Australian Clinical Trials website](#)

The anticancer drug(s) in this protocol may have been included in the ADDIKD guideline. Dose recommendations in kidney dysfunction have yet to be updated to align with the ADDIKD guideline. Recommendations will be updated once the individual protocol has been evaluated by the reference committee. For further information refer to the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\) calculator](#).

International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADIKD)

2022

[Click here](#)



Treatment schedule - Overview

Drug	Dose	Route	Day
cARBOplatin	7 AUC *	IV infusion	1

*direct measurement of renal function (e.g. by EDTA or DTPA GFR scan) is preferred whenever feasible. If estimated GFR is greater than 125 mL/min (i.e. 7 AUC dose greater than 1050mg), obtaining direct measurement rather than a calculated renal function is strongly recommended. Dose capping is strongly discouraged.

Cycles: 1

Notes:

Fertility is unlikely to be affected by 1 cycle of carboplatin however, issues including preservation (e.g. sperm banking) should be discussed with the patient prior to initiating chemotherapy.

Drug status: Carboplatin is on the [PBS general schedule](#)

Cost: ~ \$110 per cycle

Treatment schedule - Detail

The supportive therapies (e.g. antiemetics, premedications, etc.), infusion times, diluents, volumes and routes of administration, if included, are listed as defaults. They may vary between institutions and can be substituted to reflect individual institutional policy.

Antiemetics if included in the treatment schedule are based upon recommendations from national and international guidelines. These are **defaults only** and may be substituted to reflect individual institutional policy. [Select here for recommended doses of alternative antiemetics.](#)

Day 1

Netupitant	300 mg (PO)	60 minutes before chemotherapy (fixed dose preparation with palonosetron)
Palonosetron	0.5 mg (PO)	60 minutes before chemotherapy (fixed dose preparation with netupitant)
Dexamethasone	8 mg (PO)	60 minutes before chemotherapy
cARBOplatin	7 AUC (IV infusion)	in 500 mL glucose 5% over 30 to 60 minutes. (Direct measurement of renal function (e.g. by EDTA or DTPA GFR scan) is preferred whenever feasible. If estimated GFR is greater than 125 mL/min (i.e. 7 AUC dose greater than 1050mg), obtaining direct measurement rather than a calculated renal function is strongly recommended. Dose capping is strongly discouraged.)

Day 2 and 3

Dexamethasone	8 mg (PO)	ONCE a day ONCE a day (or in divided doses) with or after food. Note: dexamethasone doses on day 2 and 3 may not be required and may be reduced or omitted at the clinicians discretion *
---------------	-----------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

* [Link to ID 7 Prevention of antineoplastic induced nausea and vomiting](#)

Cycles: 1

Indications and patient population

- Adjuvant high risk Stage 1 seminoma after orchiectomy
 - Alternative to surveillance or para-aortic +/- pelvic radiation

Clinical information

Venous access required	IV cannula (IVC) or central venous access device (CVAD) is required to administer this treatment. Read more about central venous access device line selection
Hypersensitivity/infusion related reaction	High risk with carboplatin. Hypersensitivity risk increases with number of cycles of carboplatin. Rechallenge with carboplatin after hypersensitivity carries a high risk of anaphylaxis, and where clinically indicated, should be undertaken with a desensitisation protocol with appropriate supports in place. Refer to local institutional policy. Read more about Hypersensitivity reaction

Emetogenicity MODERATE	<p>Suggested default antiemetics have been added to the treatment schedule, and may be substituted to reflect institutional policy.</p> <p>Carboplatin AUC ≥ 4 is classified by MASCC/ESMO Antiemetic Guidelines 2016 and ASCO Antiemetic Guidelines 2017 as having moderate emetogenicity.</p> <p>However, a NK1 receptor antagonist and a 5HT₃ receptor antagonist in combination with dexamethasone are available on the PBS for primary prophylaxis of carboplatin induced nausea and vomiting.</p> <p>Ensure that patients also have sufficient antiemetics for breakthrough emesis:</p> <p>Metoclopramide 10 mg three times a day when necessary (maximum of 30 mg/24 hours, up to 5 days) OR</p> <p>Prochlorperazine 10 mg PO every 6 hours when necessary.</p> <p>Read more about preventing anti-cancer therapy induced nausea and vomiting</p>
Blood tests	<p>FBC, EUC and LFTs at baseline. Repeat day 28 post treatment to ensure recovery.</p>
Hepatitis B screening and prophylaxis	<p>Routine screening for HBsAg and anti-HBc is recommended prior to initiation of treatment. Prophylaxis should be determined according to individual institutional policy.</p> <p>Read more about hepatitis B screening and prophylaxis in cancer patients requiring cytotoxic and/or immunosuppressive therapy</p>
Vaccinations	<p>Live vaccines are contraindicated in cancer patients receiving immunosuppressive therapy and/or who have poorly controlled malignant disease.</p> <p>Refer to the recommended schedule of vaccination for immunocompromised patients, as outlined in the Australian Immunisation Handbook.</p> <p>Read more about COVID-19 vaccines and cancer.</p>
Fertility and fathering a child	<p>Cancer treatment can have harmful effects on fertility and this should be discussed with all patients of reproductive potential prior to commencing treatment. It is important that all patients of reproductive potential use effective contraception whilst on therapy and after treatment finishes. Effective contraception methods and contraception timeframe should be discussed with all patients of reproductive potential.</p> <p>Read more about the effect of cancer treatment on fertility</p>

Dose modifications

Evidence for dose modifications is limited, and the recommendations made on eviQ are intended as a guide only. They are generally conservative with an emphasis on safety. Any dose modification should be based on clinical judgement, and the individual patient's situation including but not limited to treatment intent (curative vs palliative), the anti-cancer regimen (single versus combination therapy versus chemotherapy versus immunotherapy), biology of the cancer (site, size, mutations, metastases), other treatment related side effects, additional co-morbidities, performance status and patient preferences. Suggested dose modifications are based on clinical trial findings, product information, published guidelines and reference committee consensus. The dose reduction applies to each individual dose and not to the total number of days or duration of treatment cycle unless stated otherwise. Non-haematological gradings are based on [Common Terminology Criteria for Adverse Events \(CTCAE\)](#) unless otherwise specified. Renal and hepatic dose modifications have been standardised where possible. For more information see dosing considerations & disclaimer.

The dose recommendations in kidney dysfunction (i.e. renal impairment) displayed may not reflect those in the ADDIKD guideline and have been included for historical reference only. Recommendations will be updated once the individual protocol has been evaluated by the reference committee, with this version of the protocol then being archived. Clinicians are expected to refer to the ADDIKD guideline prior to prescribing in kidney dysfunction.

[International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction \(ADDIKD\)](#).

For dosing carboplatin, ADDIKD recommends that:

- Directly measured glomerular filtration rate (mGFR) is the preferred kidney function value in the Calvert formula, especially where estimated kidney function may be unreliable for accurate therapeutic dosing.
- Where mGFR is unavailable, eGFR adjusted to an individual's body surface area (BSA-adjusted eGFR) is a suitable alternative for use in the Calvert formula.
- Kidney function should not be capped at 125 mL/min for use in the Calvert formula.

- Recalculation of carboplatin doses at each cycle is unnecessary, except when baseline kidney function (e.g., eGFR) alters by > 20% or when there is a change in the clinical status of the patient.

For further information refer the [eviQ Factsheet](#) around carboplatin dosing and the carboplatin drug monograph within the ADDIKD guideline. To assist with calculations, use the eviQ [Estimated Glomerular Filtration Rate \(eGFR\)](#) and [carboplatin dose calculators](#).

Note: Delay and dose reductions are not recommended as the efficacy of this treatment may be greatly compromised - clinicians are advised to consult with a medical oncologist from a tertiary treatment centre with high volume experience in testicular cancer if dose delay or reduction due to toxicities or renal or hepatic dysfunction is being contemplated; an alternative treatment regimen may need to be considered.

Interactions

Drug interactions in eviQ protocols are under review and being updated to align with current literature. Further site-wide updates and changes will occur in due course. [References & Disclaimer](#)

The drug interactions shown below are not an exhaustive list. For a more comprehensive list and for detailed information on specific drug interactions and clinical management, please refer to the specific drug product information and the following key resources:

- [MIMS - interactions tab](#) (includes link to a CYP-450 table) (login required)
- [Australian Medicines Handbook \(AMH\) – interactions tab](#) (login required)
- [Micromedex Drug Interactions](#) (login required)
- [Cancer Drug Interactions](#)
- [Cytochrome P450 Drug Interactions](#)

Carboplatin

	Interaction	Clinical management
Nephrotoxic drugs (e.g. aminoglycosides, amphotericin, contrast dye, frusemide, NSAIDs)	Additive nephrotoxicity	Avoid combination or monitor kidney function closely
Ototoxic drugs (e.g. aminoglycosides, frusemide, NSAIDs)	Additive ototoxicity	Avoid combination or perform regular audiometric testing
Paclitaxel	Administration schedule may influence the development of myelosuppression	Minimise toxicity by administering paclitaxel first in regimens using the combination

NK-1 antagonist e.g. aprepitant, fosaprepitant, netupitant		
	Interaction	Clinical management
Dexamethasone	Increased effects/toxicity of dexamethasone due to inhibition of its metabolism via CYP3A4	Reduce dose of antiemetic dexamethasone by approximately 50% when adding a NK-1 antagonist. For protocols that already recommend a NK-1 antagonist, the dose reduction of antiemetic dexamethasone has already been taken into account. If dexamethasone is part of the chemotherapy protocol , dose reduction as per the product information is not routinely recommended in clinical practice and no additional dexamethasone is required for antiemetic cover.
Warfarin	Reduced anticoagulant efficacy of warfarin due to increased clearance (aprepitant induces CYP2C9). *Note interaction only applicable to aprepitant/ fosaprepitant	INR should be monitored in the 2 week period, particularly at 7 to 10 days following the administration of aprepitant/ fosaprepitant
Combined oral contraceptive	Reduced contraceptive efficacy due to increased clearance. *Note interaction only applicable to aprepitant/ fosaprepitant	Alternative non-hormonal methods should be used during and for 1 month after stopping aprepitant/ fosaprepitant
CYP3A4 inducers (e.g. carbamazepine, phenytoin, phenobarbitone, rifampicin, St John's wort etc.)	Reduced efficacy of NK-1 antagonist possible due to increased clearance	Avoid combination or monitor for decreased antiemetic effect. Consider using an alternative antiemetic regimen
CYP3A4 inhibitors (e.g. azole antifungals, clarithromycin, erythromycin, grapefruit juice, ritonavir etc.)	Increased toxicity of NK-1 antagonist possible due to reduced clearance	Avoid combination or monitor for increased adverse effects of NK-1 antagonist (e.g. headache, hiccups, constipation)
Drugs metabolised by CYP3A4 (e.g. etoposide, imatinib, irinotecan, midazolam, paclitaxel, vinblastine, vincristine etc.)	Increased effects/toxicity of these drugs possible due to inhibition of CYP3A4 by NK-1 antagonist	Avoid combination or monitor for increased toxicity especially with orally administered drugs

General		
	Interaction	Clinical management
Warfarin	Anti-cancer drugs may alter the anticoagulant effect of warfarin.	Monitor INR regularly and adjust warfarin dosage as appropriate; consider alternative anticoagulant.
Direct oral anticoagulants (DOACs) e.g. apixaban, rivaroxaban, dabigatran	Interaction with both CYP3A4 and P-gp inhibitors /inducers. DOAC and anti-cancer drug levels may both be altered, possibly leading to loss of efficacy or toxicity (i.e. increased bleeding).	Apixaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. If treating VTE, avoid use with strong CYP3A4 and P-gp inducers. Rivaroxaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. Dabigatran: avoid combination with strong P-gp inducers and inhibitors. If concurrent use is unavoidable, monitor closely for efficacy/toxicity of both drugs.
Digoxin	Anti-cancer drugs can damage the lining of the intestine; affecting the absorption of digoxin.	Monitor digoxin serum levels; adjust digoxin dosage as appropriate.
Antiepileptics	Both altered antiepileptic and anti-cancer drug levels may occur, possibly leading to loss of efficacy or toxicity.	Where concurrent use of an enzyme-inducing antiepileptic cannot be avoided, monitor antiepileptic serum levels for toxicity, as well as seizure frequency for efficacy; adjust dosage as appropriate. Also monitor closely for efficacy of the anti-cancer therapy.
Antiplatelet agents and NSAIDs	Increased risk of bleeding due to treatment related thrombocytopenia.	Avoid or minimise combination. If combination deemed essential, (e.g. low dose aspirin for ischaemic heart disease) monitor for signs of bleeding.
Serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs e.g. paroxetine) and serotonin noradrenaline reuptake inhibitors (SNRIs e.g. venlafaxine)	Increased risk of serotonin syndrome with concurrent use of 5-HT3 receptor antagonists (e.g. palonosetron, ondansetron, granisetron, tropisetron, dolasetron, etc.)	Avoid combination. If combination is clinically warranted, monitor for signs and symptoms of serotonin syndrome (e.g. confusion, agitation, tachycardia, hyperreflexia). For more information link to TGA Medicines Safety Update
Vaccines	Diminished response to vaccines and increased risk of infection with live vaccines.	Live vaccines (e.g. BCG, MMR, zoster and varicella) are contraindicated in patients on immunosuppressive therapy. Use with caution in patients on non-immunosuppressive therapy. For more information; refer to the recommended schedule of vaccination for cancer patients, as outlined in the Australian Immunisation Handbook

Administration

eviQ provides safe and effective instructions on how to administer cancer treatments. However, eviQ does not provide every treatment delivery option, and is unable to provide a comprehensive list of cancer treatment agents and their required IV line giving set/filter. There may be alternative methods of treatment administration, and alternative supportive treatments that are also appropriate. Please refer to the individual

Day 1

Approximate treatment time: 90 minutes

[Safe handling and waste management](#)

[Safe administration](#)

[General patient assessment](#) prior to each day of treatment.

Any toxicity grade 2 or greater may require delay of treatment and review by medical officer before commencing treatment.

Prime IV line(s).

Insert IV cannula or access [TIVAD](#) or [CVAD](#).

Pre treatment medication

Verify antiemetics taken or administer as prescribed.

Verify dexamethasone taken or administer as prescribed.

🕒 Chemotherapy - Time out

Carboplatin

Administer carboplatin (irritant):

- via IV infusion over 30 to 60 minutes
- observe for hypersensitivity reactions
- flush with ~100 mL of sodium chloride 0.9%
- hypersensitivity risk increases with number of cycles administered.

Stop infusion at first sign of reaction:

- if symptoms are mild and resolve when infusion is stopped, consider recommencing infusion after review by medical officer at a slower rate
- for severe reactions seek medical assistance immediately and do not restart infusion.

Remove IV cannula and/or deaccess [TIVAD](#) or [CVAD](#).

Continue [safe handling](#) precautions until 7 days after completion of drug(s)

Discharge Information

Antiemetics

- Antiemetics as prescribed.

Patient information

- Ensure patient receives patient information sheet.

Side effects

The side effects listed below are not a complete list of all possible side effects for this treatment. Side effects are categorised into the approximate onset of presentation and should only be used as a guide.

Immediate (onset hours to days)	
Hypersensitivity reaction	Anaphylaxis and infusion related reactions can occur with this treatment. Read more about hypersensitivity reaction
Nausea and vomiting	Read more about prevention of treatment induced nausea and vomiting
Taste and smell alteration	Read more about taste and smell changes
Early (onset days to weeks)	
Neutropenia	Abnormally low levels of neutrophils in the blood. This increases the risk of infection. Any fever or suspicion of infection should be investigated immediately and managed aggressively. Read more about immediate management of neutropenic fever
Thrombocytopenia	A reduction in the normal levels of functional platelets, increasing the risk of abnormal bleeding. Read more about thrombocytopenia
Oral mucositis	Erythematous and ulcerative lesions of the gastrointestinal tract (GIT). It commonly develops following chemotherapy, radiation therapy to the head, neck or oesophagus, and high dose chemotherapy followed by a blood and marrow transplant (BMT). Read more about oral mucositis
Fatigue	Read more about fatigue
Late (onset weeks to months)	
Anaemia	Abnormally low levels of red blood cells (RBCs) or haemoglobin in the blood. Read more about anaemia
Alopecia - partial	Hair thinning and/or patchy hair loss. Patients can also experience mild to moderate discomfort of the hair follicles, and rarely pain as the hair is falling out. Read more about alopecia and scalp cooling

Evidence

An EORTC/MRC study randomised 1477 stage 1 seminoma patients to either a single dose of carboplatin (AUC 7) or radiation therapy (para-aortic strip or dog-leg). 904 and 573 patients received radiation therapy and carboplatin, respectively.


With a median follow-up of 6.5 years, relapse-free survival rates for radiation therapy and carboplatin were similar (96% vs 94.7% at 5 years, respectively; hazard ratio 1.25; p=0.37).¹ Therefore confirming the non-inferiority of single dose carboplatin versus radiation therapy in terms of relapse free rate. A statistically significant reduction medium term risk of second germ cell tumours with carboplatin was also demonstrated (Carboplatin n=2; RT n=15; hazard ratio, 0.22; p=0.03). This update did not contain any new toxicity data.

In the original report patients given carboplatin were less lethargic and less likely to take time off work than those given radiation therapy. New, second primary testicular germ-cell tumours were reported in ten patients allocated irradiation (all after para-aortic strip field) and two allocated carboplatin. One seminoma-related death occurred after radiation therapy and none after carboplatin. Preliminary evidence indicated a reduced incidence of cancer in the contra-lateral testis.²

Efficacy

New, second primary testicular germ-cell tumours were reported in ten patients allocated irradiation (all after para-aortic strip field) and two allocated carboplatin. One seminoma-related death occurred after radiation therapy and none after carboplatin.²

Relapse-free rate allocated by treatment:¹



@ JCO 2011

Toxicity

There were significantly higher rates of grade 3-4 thrombocytopenia in the patients receiving carboplatin 17 (4%) vs 0; $p < 0.0001$ and significantly more dyspepsia in those receiving radiation therapy 127 (17%) vs 40 (8%); $p < 0.0001$. Patients given carboplatin were less lethargic and less likely to take time off work than those given radiation therapy. New, second primary testicular germ-cell tumours were reported in ten patients allocated irradiation (all after para-aortic strip field) and two allocated carboplatin. One seminoma-related death occurred after radiation therapy and none after carboplatin.²

References

- 1 Oliver, R. T., Mead, G.M., Rustin, G.J. 2011. "Randomized trial of carboplatin versus radiotherapy for stage I seminoma: mature results on relapse and contralateral testis cancer rates in MRC TE19/EORTC 30982 study (ISRCTN27163214)". *J Clin Oncol.* 29 (8): 957-62.
- 2 Oliver, R. T., M. D. Mason, G. M. Mead, et al. 2005. "Radiotherapy versus single-dose carboplatin in adjuvant treatment of stage I seminoma: a randomised trial." *Lancet.* 366(9482):293-300.

History

Version 4

Date	Summary of changes
09/05/2007	Patient sheet updated.
06/07/2007	Minor changes with annual review.
19/01/2010	Added discussion regarding one versus two cycles of carboplatin.
15/02/2010	Review, new dose modifications and transferred to eviQ.
17/05/2010	Decision by reference committee to keep the preferred number of cycles as 1.
21/05/2011	New format to allow for export of protocol information. Protocol version number changed to V.2. Antiemetics and premedications added to the treatment schedule. Additional Clinical Information, Key Prescribing table and Key Administration table combined into new section titled Clinical Considerations.

Date	Summary of changes
	Drug specific information placed behind the drug name link. Addition of blood tests: repeat bloods Day 28 post treatment to ensure recovery.
09/09/2011	Infusion fluid for carboplatin changed from sodium chloride 0.9% to glucose 5% because of longer stability.
02/12/2011	Peripheral neuropathy.
16/01/2012	PHC view updated.
18/04/2012	Palonosetron added as the preferred 5HT ₃ antagonist for moderate emetogenicity.
30/11/2012	Reviewed at Reference Committee meeting. Evidence update supporting 1 cycle only. Protocol updated.
09/05/2014	Reviewed at Medical Oncology Reference Committee Meeting. Note about measured GFR added. PHC view removed. Review 2 years.
31/03/2017	Protocol discussed and decided to have a 5 year review period. Next due for review in 2019.
31/05/2017	Transferred to new eviQ website. Version number changed to V.3. Antiemetic change: A NK1 receptor antagonist and a 5HT ₃ receptor antagonist in combination with dexamethasone has been added as available on the PBS for primary prophylaxis of carboplatin induced nausea and vomiting.
10/05/2018	Dose modifications updated to delay or dose reduction not recommended wording consistent with all testicular protocols and as single dose treatment.
17/01/2019	Carboplatin AUC ≥ 4 changed from highly to moderately emetogenic as per MASCC/ESMO and ASCO guidelines and medical oncology reference committee consensus. Dexamethasone day 4 dose removed. NK1 receptor antagonist unchanged. Treatment detail and clinical information updated to reflect the change. Version number changed to V.4
25/03/2019	Protocol reviewed at Medical Oncology Reference Committee meeting on 15/03/2019. ANZUP surveillance recommendations added as related page. Next review in 5 years.
31/10/2019	Note regarding renal function and dose capping under treatment schedule updated.

The information contained in this protocol is based on the highest level of available evidence and consensus of the eviQ reference committee regarding their views of currently accepted approaches to treatment. Any clinician (medical oncologist, haematologist, radiation oncologist, medical physicist, radiation therapist, pharmacist or nurse) seeking to apply or consult this protocol is expected to use independent clinical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use is subject to eviQ's disclaimer available at www.eviq.org.au

First approved: 14 September 2005

Last reviewed: 15 March 2019

Review due: 30 June 2024

The currency of this information is guaranteed only up until the date of printing, for any updates please check:

<https://www.eviq.org.au/p/325>

19 Jun 2023

Patient information - Testicular cancer adjuvant - Carboplatin

Patient's name:

Your treatment

The treatment schedule below explains how the drug for this treatment is given.

Carboplatin			
This treatment is given once.			
Day	Treatment	How it is given	How long it takes
1	Carboplatin (<i>carb-o-PLAT-in</i>)	By a drip into a vein	About 1.5 hours

When to get help

Anticancer drugs (drugs used to treat cancer) can sometimes cause serious problems. It is important to get medical help immediately if you become unwell.

 <p>IMMEDIATELY go to your nearest hospital Emergency Department, or contact your doctor or nurse if you have any of the following at any time:</p>	Emergency contact details Ask your doctor or nurse from your treating team who to contact if you have a problem
<ul style="list-style-type: none">• a temperature of 38°C or higher• chills, sweats, shivers or shakes• shortness of breath• uncontrolled vomiting or diarrhoea• pain, tingling or discomfort in your chest or arms• you become unwell.	Daytime:..... Night/weekend:..... Other instructions:.....

During your treatment immediately tell the doctor or nurse looking after you if you get any of the following problems:

- leaking from the area where the drugs are being given
- pain, stinging, swelling or redness in the area where the drugs are being given or at any injection sites
- a skin rash, itching, feeling short of breath, wheezing, fever, shivers, or feeling dizzy or unwell in any way (allergic reaction).

Other information about your treatment

Treatment delays

There may be times when your treatment is delayed. This can happen if your doctor thinks you are likely to have severe side effects, if you get severe side effects, if your blood counts are affected and causing delays in treatment, or if you are finding it hard to cope with the treatment. Your doctor will explain if you need any delays to your treatment and the reason why.

You will need to have a blood test before you start treatment. Your doctor or nurse will tell you when to have the blood test.

Other medications given during this treatment

- **Anti-sickness (anti-nausea) medication:** you may be given some anti-sickness medication. Make sure you take this medication as your doctor or nurse tells you, even if you don't feel sick. This can help to prevent the sickness starting.

Side effects

Cancer treatments can cause damage to normal cells in your body, which can cause side effects. Everyone gets different side effects, and some people will have more problems than others.

The table below shows some of the side effects you may get with this treatment. You are unlikely to get all of those listed and you may also get some side effects that have not been listed.

Tell your doctor or nurse about any side effects that worry you. Follow the instructions below and those given to you by your doctor or nurse.

Immediate (onset hours to days)	
Allergic reaction	<ul style="list-style-type: none">• Allergic reactions are uncommon but can be life threatening.• If you feel unwell during the infusion or shortly after it, or:<ul style="list-style-type: none">◦ get a fever, shivers or shakes◦ feel dizzy, faint, confused or anxious◦ start wheezing or have difficulty breathing◦ have a rash, itch or redness of the face <p><u>While you are in hospital:</u> Tell your doctor or nurse immediately.</p> <p><u>After you leave:</u> Contact your doctor or nurse immediately, or go to the nearest hospital Emergency Department.</p>
Nausea and vomiting	<ul style="list-style-type: none">• You may feel sick (nausea) or be sick (vomit).• Take your anti-sickness medication as directed even if you don't feel sick.• Drink plenty of fluids (unless you are fluid restricted).• Eat small meals more frequently.• Try food that does not require much preparation.• Try bland foods like dry biscuits or toast.• Gentle exercise may help with nausea.• Ask your doctor or nurse for eviQ patient information - Nausea and vomiting during cancer treatment.• Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have uncontrolled vomiting or feel dizzy or light-headed.
Taste and smell changes	<ul style="list-style-type: none">• You may find that food loses its taste or tastes different.• These changes are likely to go away with time.• Do your mouth care regularly.• Chew on sugar-free gum or eat sugar-free mints.• Add flavour to your food with sauces and herbs.• Ask your doctor or nurse for eviQ patient information - Taste and smell changes during cancer treatment.

Early (onset days to weeks)

<p>Infection risk (neutropenia)</p>	<ul style="list-style-type: none"> • This treatment lowers the amount of white blood cells in your body. The type of white blood cells that help to fight infection are called neutrophils. Having low level of neutrophils is called neutropenia. If you have neutropenia, you are at greater risk of getting an infection. It also means that your body can't fight infections as well as usual. This is a serious side effect, and can be life threatening. • Wash your hands often. • Keep a thermometer at home and take your temperature regularly, and if you feel unwell. • Do your mouth care regularly. • Inspect your central line site (if you have one) daily for any redness, pus or swelling. • Limit contact with people who are sick. • Learn how to recognise the signs of infection. • Ask your doctor or nurse for eviQ patient information - Infection during cancer treatment. • Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the following signs or symptoms: <ul style="list-style-type: none"> ◦ a temperature of 38°C or higher ◦ chills, shivers, sweats or shakes ◦ a sore throat or cough ◦ uncontrolled diarrhoea ◦ shortness of breath ◦ a fast heartbeat ◦ become unwell even without a temperature.
<p>Low platelets (thrombocytopenia)</p>	<ul style="list-style-type: none"> • This treatment lowers the amount of platelets in your blood. Platelets help your blood to clot. When they are low, you are at an increased risk of bleeding and bruising. • Try not to bruise or cut yourself. • Avoid contact sport or vigorous exercise. • Clear your nose by blowing gently. • Avoid constipation. • Brush your teeth with a soft toothbrush. • Don't take aspirin, ibuprofen or other similar anti-inflammatory medications unless your doctor tells you to. • Tell your doctor or nurse if you have any bruising or bleeding. • Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if you have any uncontrolled bleeding.
<p>Mouth pain and soreness (mucositis)</p>	<ul style="list-style-type: none"> • You may have: <ul style="list-style-type: none"> ◦ bleeding gums ◦ mouth ulcers ◦ a white coating on your tongue ◦ pain in the mouth or throat ◦ difficulty eating or swallowing. • Avoid spicy, acidic or crunchy foods and very hot or cold food and drinks. • Try bland and soft foods. • Brush your teeth gently with a soft toothbrush after each meal and at bedtime. If you normally floss continue to do so. • Rinse your mouth after you eat and brush your teeth, using either: <ul style="list-style-type: none"> ◦ 1/4 teaspoon of salt in 1 cup of warm water, or ◦ 1/4 teaspoon of bicarbonate of soda in 1 cup of warm water • Ask your doctor or nurse for eviQ patient information - Mouth problems during cancer treatment. • Tell your doctor or nurse if you get any of the symptoms listed above.

Tiredness and lack of energy (fatigue)	<ul style="list-style-type: none"> You may feel very tired, have no energy, sleep a lot, and not be able to do normal activities or things you enjoy. Do not drive or operate machinery if you are feeling tired. Nap for short periods (only 1 hour at a time) Prioritise your tasks to ensure the best use of your energy. Eat a well balanced diet and drink plenty of fluids (unless you are fluid restricted). Try some gentle exercise daily. Allow your friends and family to help. Tell your doctor or nurse if you get any of the symptoms listed above.
Late (onset weeks to months)	
Low red blood cells (anaemia)	<ul style="list-style-type: none"> You may feel dizzy, light-headed, tired and appear more pale than usual. Tell your doctor or nurse if you have any of these signs or symptoms. You might need a blood transfusion. Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have any chest pain, trouble breathing, or feel like your heart is racing.
Hair thinning	<ul style="list-style-type: none"> Your hair may become dry and may break easily. You may lose some of your hair. Use a gentle shampoo and a soft hairbrush. Take care with hair products like hairspray, hair dye, bleaches and perms. Protect your scalp from the cold with a hat or scarf. Protect your scalp from the sun with a hat and sunscreen of SPF 50 or higher. Ask your doctor or nurse about the Look Good Feel Better program (www.lgfb.org.au)

General advice for people having cancer treatment

Chemotherapy safety

- Learn how to keep you and your family safe while you are having anticancer drugs.
- See our patient information sheet - [Chemotherapy safety at home](#).

Blood clot risk

- Cancer and anticancer drugs can increase the risk of a blood clot (thrombosis).
- Tell your doctor if you have a family history of blood clots.
- A blood clot can cause pain, redness, swelling in your arms or legs, shortness of breath or chest pain.
- If you have any of these symptoms go to your nearest hospital Emergency Department.

Medications and vaccinations

- Before you start treatment, tell your doctor about any medications you are taking, including vitamins or herbal supplements.
- Don't stop or start any medications during treatment without talking to your doctor and pharmacist first.
- Paracetamol is safe to take if you have a headache or other mild aches and pains. It is recommended that you avoid taking aspirin, ibuprofen and other anti-inflammatory type medications for pain while you are having treatment. However, if these medications have been prescribed by your doctor, do not stop taking them without speaking with your doctor.
- Vaccinations such as flu and tetanus vaccines are safe to receive while having treatment. Do not have any live vaccines during your treatment or for 6 months after it finishes. If you are unsure, check with your doctor before you have any vaccinations.
- People you live with should be fully vaccinated, including having live vaccines according to the current vaccination schedule. Extra care needs to be taken with hand washing and careful disposal of soiled nappies for infants who have recently received the rotavirus vaccine.

Other medical and dental treatment

- If you go to hospital or any other medical appointment (including dental appointments), always tell the person treating you that you are receiving anticancer drugs.
- Before you have any dental treatment, talk to your doctor.

Diet

- While you are receiving this treatment it is important that you try to maintain a healthy diet.
- Speak to your doctor or nurse about whether drinking alcohol is safe with your treatment.
- If you have any concerns about recent weight loss or weight gain or questions about your diet, ask to speak to a dietitian.

Fertility

- Some cancer treatments can reduce your fertility. This can make it difficult or impossible to father a child.
- Talk to your doctor or nurse before you start any treatment. Depending on your situation there may be fertility sparing options available to you and/or your partner, discuss these with your doctor or nurse.

Fathering a child

- Some cancer treatments can be dangerous to unborn babies. Talk to your doctor or nurse if you think there is any chance that your partner could be pregnant.
- Do not try to father a child during this treatment. Contraception should be used during treatment and after stopping treatment. Ask your doctor or nurse about what type of contraception you should use.
- If you are planning fatherhood after completing this treatment, talk to your doctor. Some doctors advise waiting between 6 months and 2 years after treatment.

Sex life and sexuality

- The desire to have sex may decrease as a result of this treatment or its side effects.
- Your emotions and the way you feel about yourself may also be affected by this treatment.
- It may help to discuss your concerns with your partner and doctor or nurse.

Quitting smoking

- It is never too late to quit smoking. Quitting smoking is one of the best things you can do to help your treatment work better.
- There are many effective tools to improve your chances of quitting.
- Talk to your treating team for more information and referral to a smoking cessation support service.

Staying active

- Research shows that exercise, no matter how small, has many benefits for people during and after cancer treatment.
- Talk to your doctor before starting an exercise program. Your doctor can advise whether you need a modified exercise program.

For more information about cancer treatment, side effects and side effect management see our [Patient and carers](#) section.

Where to get more information

Telephone support

- Call Cancer Council on 13 11 20 for cancer information and support

Testicular cancer information

- Healthy Male Andrology Australia – healthymale.org.au
- Livestrong – livestrong.org

General cancer information and support

- Australian Rare Cancer (ARC) Portal – arcportal.org.au/
- Beyondblue – beyondblue.org.au
- Cancer Australia – canceraustralia.gov.au
- Cancer Council Australia – cancer.org.au
- Cancer Voices Australia – cancervoicesaustralia.org
- CanTeen – canteen.org.au
- Carers Australia – carersaustralia.com.au
- CHILL Cancer related hair loss - scalpcooling.org
- eviQ Cancer Treatments Online – eviQ.org.au
- LGBTQI+ People and Cancer - cancercouncil.com.au/cancer-information/lgbtqi
- Look Good Feel Better – lgfb.org.au
- Patient Information – patients.cancer.nsw.gov.au

