

# Colorectal adjuvant de Gramont (fluorouracil and leucovorin) SUPERSEDED

ID: 75 v.5 **Superseded** Essential Medicine List

This protocol has been superseded because it is not commonly used in clinical practice and [Colorectal adjuvant de Gramont modified \(ID 76\)](#) is the preferred regimen.

## ⚠ Fluoropyrimidine overdose or overexposure:

Fluoropyrimidine overdose or overexposure may result in severe or life-threatening toxicity. An antidote is available and is highly effective if given within 96 hours. Read more about [fluoropyrimidine overdose or overexposure](#).

Check for clinical trials in this patient group. Link to [Australian Clinical Trials](#) website

The anticancer drug(s) in this protocol may have been included in the ADDIKD guideline. Dose recommendations in kidney dysfunction have yet to be updated to align with the ADDIKD guideline. Recommendations will be updated once the individual protocol has been evaluated by the reference committee. For further information refer to the ADDIKD guideline. To assist with calculations, use the [eviQ Estimated Glomerular Filtration Rate \(eGFR\) calculator](#).

## International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADDIKD)

2022

[Click here](#)



### Related pages:

- [Colorectal adjuvant de Gramont \(modified\) \(fluorouracil and leucovorin\)](#)

## Treatment schedule - Overview

### Cycle 1 to 12

Drug	Dose	Route	Day
Calcium folinate (Leucovorin)	200 mg/m <sup>2</sup> *	IV infusion	1 and 2
Fluorouracil	400 mg/m <sup>2</sup>	IV	1 and 2
Fluorouracil	600 mg/m <sup>2</sup>	CIV via pump over 22 hours	1 and 2

\* This regimen uses calcium folinate (Leucovorin<sup>®</sup>) at a dose of 200 mg/m<sup>2</sup>, as defined in the original clinical trials. This dose is administered over a time period of 2 hours which increases the clinic time from 30 minutes to 2.5 hours. A discussion regarding the effect of dosing on outcome can be found in the calcium folinate dose document.

**Frequency:** 14 days

**Cycles:** 12

**Drug status:** All drugs in this protocol are on the [PBS general schedule](#)

**Cost:** ~ \$230 per cycle

## Treatment schedule - Detail

The supportive therapies (e.g. antiemetics, premedications, etc.), infusion times, diluents, volumes and routes of administration, if included, are listed as defaults. They may vary between institutions and can be substituted to reflect individual institutional policy.

Antiemetics if included in the treatment schedule are based upon recommendations from national and international guidelines. These are **defaults only** and may be substituted to reflect individual institutional policy. Select here for recommended doses of alternative antiemetics.

## Cycle 1 to 12

Day 1 and 2		
Metoclopramide	10 mg (PO)	one tablet when necessary (maximum of 30 mg/24 hours, up to 5 days)
Calcium folinate (Leucovorin)	200 mg/m <sup>2</sup> (IV infusion)	in 250 mL sodium chloride 0.9% over 2 hours
Fluorouracil	400 mg/m <sup>2</sup> (IV)	over 3 to 5 minutes
Fluorouracil	600 mg/m <sup>2</sup> (CIV)	via ambulatory infusion pump over 22 hours

This regimen uses calcium folinate (Leucovorin<sup>®</sup>) at a dose of 200 mg/m<sup>2</sup>, as defined in the original clinical trials. This dose is administered over a time period of 2 hours which increases the clinic time from 30 minutes to 2.5 hours. A discussion regarding the effect of dosing on outcome can be found in the calcium folinate dose document.

**Frequency:** 14 days

**Cycles:** 12

## Indications and patient population

- Adjuvant colorectal cancer.

## Clinical information

<b>Safety alert fluoropyrimidines</b>	Fluoropyrimidines can be administered by different routes and schedules with each method having associated increased risk of certain side effects. Fluoropyrimidine overdose or overexposure is a rare but potentially life threatening side effect of this drug class and can occur by any route of administration. An antidote is available and highly effective if given within 96 hours.  Read more about the <a href="#">medication safety alert for infusional fluorouracil and fluoropyrimidine overdose or overexposure</a>
<b>Venous access</b>	Central venous access device (CVAD) is required to administer this treatment.  Read more about <a href="#">central venous access device line selection</a>
<b>Emetogenicity LOW</b>	Suggested default antiemetics have been added to the treatment schedule, and may be substituted to reflect institutional policy.  Ensure that patients also have sufficient antiemetics for breakthrough emesis: Metoclopramide 10 mg three times a day when necessary (maximum of 30 mg/24 hours, up to 5 days) OR Prochlorperazine 10 mg PO every 6 hours when necessary.  Read more about <a href="#">preventing anti-cancer therapy induced nausea and vomiting</a>
<b>Cardiac toxicity</b>	Cardiac toxicity is a serious complication that can occur during treatment with fluorouracil. Patients treated with fluorouracil, especially those with a prior history of cardiac disease or other risk factors, should be carefully monitored during therapy.  Read more about <a href="#">cardiac toxicity associated with anti-cancer drugs</a>

<b>Dihydropyrimidine dehydrogenase (DPD) enzyme deficiency</b>	Rare, life-threatening toxicities such as mucositis, neutropenia, neurotoxicity and diarrhoea have been reported following administration of fluoropyrimidines (e.g. fluorouracil and capecitabine). Severe unexplained toxicities require investigation prior to continuing with treatment. Testing for DPD enzyme deficiency is available in Australia but not currently reimbursed.  Read more about <a href="#">dihydropyrimidine dehydrogenase (DPD) enzyme deficiency</a>
<b>Severe enteropathy associated with fluoropyrimidine</b>	Severe enteropathy has been reported among patients with stage II/III colon cancer treated with fluoropyrimidine chemotherapy with or without oxaliplatin. Patients treated with fluoropyrimidine should be closely monitored for diarrhoea and aggressively managed.  Read more about <a href="#">severe enteropathy associated with fluorouracil in colorectal cancer</a>
<b>Diarrhoea</b>	Antidiarrhoeals (e.g. loperamide) are usually prescribed with this treatment.  Read more about <a href="#">treatment induced diarrhoea</a>
<b>Growth factor support</b>	G-CSF (short or long-acting) is available on the PBS for chemotherapy induced neutropenia depending on clinical indication and/or febrile neutropenia risk.  Access the <a href="#">PBS website</a>
<b>Blood tests</b>	FBC, EUC and LFTs at baseline and prior to each cycle. INR as clinically indicated.
<b>Hepatitis B screening and prophylaxis</b>	Routine screening for HBsAg and anti-HBc is recommended prior to initiation of treatment. Prophylaxis should be determined according to individual institutional policy.  Read more about <a href="#">hepatitis B screening and prophylaxis in cancer patients requiring cytotoxic and/or immunosuppressive therapy</a>
<b>Vaccinations</b>	Live vaccines are contraindicated in cancer patients receiving immunosuppressive therapy and/or who have poorly controlled malignant disease.  Refer to the recommended schedule of vaccination for immunocompromised patients, as outlined in the <a href="#">Australian Immunisation Handbook</a> .  Read more about <a href="#">COVID-19 vaccines and cancer</a> .
<b>Fertility, pregnancy and lactation</b>	Cancer treatment can have harmful effects on fertility and this should be discussed with all patients of reproductive potential prior to commencing treatment. There is a risk of foetal harm in pregnant women. A pregnancy test should be considered prior to initiating treatment in females of reproductive potential if sexually active. It is important that all patients of reproductive potential use effective contraception whilst on therapy and after treatment finishes. Effective contraception methods and adequate contraception timeframe should be discussed with all patients of reproductive potential. Possibility of infant risk should be discussed with breastfeeding patients.  Read more about the <a href="#">effect of cancer treatment on fertility</a>

## Dose modifications

*Evidence for dose modifications is limited, and the recommendations made on eviQ are intended as a guide only. They are generally conservative with an emphasis on safety. Any dose modification should be based on clinical judgement, and the individual patient's situation including but not limited to treatment intent (curative vs palliative), the anti-cancer regimen (single versus combination therapy versus chemotherapy versus immunotherapy), biology of the cancer (site, size, mutations, metastases), other treatment related side effects, additional co-morbidities, performance status and patient preferences. Suggested dose modifications are based on clinical trial findings, product information, published guidelines and reference committee consensus. The dose reduction applies to each individual dose and not to the total number of days or duration of treatment cycle unless stated otherwise. Non-haematological gradings are based on [Common Terminology Criteria for Adverse Events \(CTCAE\)](#) unless otherwise specified. Renal and hepatic dose modifications have been standardised where possible. For more information see dosing considerations & disclaimer.*

The dose recommendations in kidney dysfunction (i.e. renal impairment) displayed may not reflect those in the ADDIKD guideline and have been included for historical reference only. Recommendations will be updated once the individual protocol has been evaluated by the reference committee, with this version of the protocol then being archived. Clinicians are expected to

refer to the ADDIKD guideline prior to prescribing in kidney dysfunction.

**International Consensus Guideline for Anticancer Drug Dosing in Kidney Dysfunction (ADDIKD).**

**Note:**

- All dose reductions are calculated as a percentage of the starting dose.
- The dose of calcium folinate (Leucovorin®) remains fixed at 200 mg/m<sup>2</sup> and is delayed or omitted if fluorouracil is delayed or omitted.

<b>Haematological toxicity</b>	
<b>ANC x 10<sup>9</sup>/L (pre-treatment blood test)</b>	
0.5 to less than 1.0	Delay treatment until recovery
less than 0.5	Delay treatment until recovery and consider reducing fluorouracil by 25% for subsequent cycles
Febrile neutropenia	Delay treatment until recovery and consider reducing fluorouracil by 25% for subsequent cycles
<b>Platelets x 10<sup>9</sup>/L (pre-treatment blood test)</b>	
75 to less than 100	Refer to local institutional guidelines; it is the view of the expert clinicians that treatment should continue if patient is clinically well
50 to less than 75	Delay treatment until recovery
less than 50	Delay treatment until recovery and consider reducing fluorouracil by 25% for subsequent cycles

<b>Renal impairment</b>	
<b>Creatinine clearance (mL/min)</b>	
30 to 50	Reduce fluorouracil by 25%
less than 30	Reduce fluorouracil by 50%

<b>Hepatic impairment</b>	
<b>Hepatic dysfunction</b>	
Mild	No dose modifications necessary
Moderate	Reduce fluorouracil by 25%
Severe	Reduce fluorouracil by 50%

<b>Mucositis and stomatitis</b>	
Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce fluorouracil by 25% 3 <sup>rd</sup> occurrence: Reduce fluorouracil by 50% 4 <sup>th</sup> occurrence: Omit fluorouracil
Grade 3 or Grade 4	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce fluorouracil by 50% 2 <sup>nd</sup> occurrence: Omit fluorouracil

<b>Diarrhoea</b>	
Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce fluorouracil by 25%

Diarrhoea	
	3 <sup>rd</sup> occurrence: Reduce fluorouracil by 50% 4 <sup>th</sup> occurrence: Omit fluorouracil
Grade 3 or Grade 4	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce fluorouracil by 50% 2 <sup>nd</sup> occurrence: Omit fluorouracil

Hand foot syndrome (link to Hand foot syndrome (Palmar-plantar erythrodysesthesia))	
Grade 2	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: No dose reduction 2 <sup>nd</sup> occurrence: Reduce fluorouracil 25% 3 <sup>rd</sup> occurrence: Reduce fluorouracil by 50% 4 <sup>th</sup> occurrence: Omit fluorouracil
Grade 3	Delay treatment until toxicity has resolved to Grade 1 or less and reduce the dose for subsequent cycles as follows: 1 <sup>st</sup> occurrence: Reduce fluorouracil by 50% 2 <sup>nd</sup> occurrence: Omit fluorouracil

## Interactions

Drug interactions in eviQ protocols are under review and being updated to align with current literature. Further site-wide updates and changes will occur in due course. *References & Disclaimer*

The drug interactions shown below are not an exhaustive list. For a more comprehensive list and for detailed information on specific drug interactions and clinical management, please refer to the specific drug product information and the following key resources:

- [MIMS - interactions tab](#) (includes link to a CYP-450 table) (login required)
- [Australian Medicines Handbook \(AMH\) – interactions tab](#) (login required)
- [Micromedex Drug Interactions](#) (login required)
- [Cancer Drug Interactions](#)
- [Cytochrome P450 Drug Interactions](#)

Fluorouracil		
	Interaction	Clinical management
<b>Folic acid</b>	Increased toxicity of fluorouracil due to stabilisation of its bond to thymidylate synthetase (folic acid is a precursor of folinic acid/leucovorin)	Advise patients not to take folic acid supplements (inc. multivitamins) around the time of receiving treatment with fluorouracil
<b>Metronidazole, tinidazole</b>	Increased toxicity of fluorouracil due to reduced clearance	Avoid combination or monitor for fluorouracil toxicity
<b>Warfarin and other drugs metabolised by CYP2C9 (e.g. warfarin, phenytoin etc.)</b>	Increased effect/toxicity of these drugs due to inhibition of CYP2C9 by fluorouracil resulting in reduced clearance	Avoid combination or monitor for increased effect/toxicity of these drugs (e.g. for bleeding/elevated INR with warfarin, elevated phenytoin serum levels or signs of toxicity such as ataxia, tremor etc.)
<b>Allopurinol</b>	Reduced efficacy of fluorouracil possible due to reduced conversion to the active metabolites	Avoid combination or monitor for reduced fluorouracil efficacy

General		
	Interaction	Clinical management
<b>Warfarin</b>	Anti-cancer drugs may alter the anticoagulant effect of warfarin.	Monitor INR regularly and adjust warfarin dosage as appropriate; consider alternative anticoagulant.
<b>Direct oral anticoagulants (DOACs) e.g. apixaban, rivaroxaban, dabigatran</b>	Interaction with both CYP3A4 and P-gp inhibitors /inducers.  DOAC and anti-cancer drug levels may both be altered, possibly leading to loss of efficacy or toxicity (i.e. increased bleeding).	Apixaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors. If treating VTE, avoid use with strong CYP3A4 and P-gp inducers.  Rivaroxaban: avoid concurrent use with strong CYP3A4 and P-gp inhibitors.  Dabigatran: avoid combination with strong P-gp inducers and inhibitors.  If concurrent use is unavoidable, monitor closely for efficacy/toxicity of both drugs.
<b>Digoxin</b>	Anti-cancer drugs can damage the lining of the intestine; affecting the absorption of digoxin.	Monitor digoxin serum levels; adjust digoxin dosage as appropriate.
<b>Antiepileptics</b>	Both altered antiepileptic and anti-cancer drug levels may occur, possibly leading to loss of efficacy or toxicity.	Where concurrent use of an enzyme-inducing antiepileptic cannot be avoided, monitor antiepileptic serum levels for toxicity, as well as seizure frequency for efficacy; adjust dosage as appropriate. Also monitor closely for efficacy of the anti-cancer therapy.
<b>Antiplatelet agents and NSAIDs</b>	Increased risk of bleeding due to treatment related thrombocytopenia.	Avoid or minimise combination. If combination deemed essential, (e.g. low dose aspirin for ischaemic heart disease) monitor for signs of bleeding.
<b>Serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs e.g. paroxetine) and serotonin noradrenaline reuptake inhibitors (SNRIs e.g. venlafaxine)</b>	Increased risk of serotonin syndrome with concurrent use of 5-HT3 receptor antagonists (e.g. palonosetron, ondansetron, granisetron, tropisetron, dolasetron, etc.)	Avoid combination. If combination is clinically warranted, monitor for signs and symptoms of serotonin syndrome (e.g. confusion, agitation, tachycardia, hyperreflexia). For more information link to <a href="#">TGA Medicines Safety Update</a>
<b>Vaccines</b>	Diminished response to vaccines and increased risk of infection with live vaccines.	Live vaccines (e.g. BCG, MMR, zoster and varicella) are contraindicated in patients on immunosuppressive therapy. Use with caution in patients on non-immunosuppressive therapy. For more information; refer to the recommended schedule of vaccination for cancer patients, as outlined in the <a href="#">Australian Immunisation Handbook</a>

## Administration

*eviQ provides safe and effective instructions on how to administer cancer treatments. However, eviQ does not provide every treatment delivery option, and is unable to provide a comprehensive list of cancer treatment agents and their required IV line giving set/filter. There may be alternative methods of treatment administration, and alternative supportive treatments that are also appropriate. Please refer to the individual*

## Day 1

**Approximate treatment time: 2.5 hours**

[Safe handling and waste management](#)

[Safe administration](#)

[General patient assessment](#) prior to each treatment.

Any toxicity grade 2 or greater may require dose reduction, delay or omission of treatment and review by medical officer before recommencing treatment.

Prime IV line(s).

Access [TIVAD](#) or [CVAD](#).

### Pre treatment medication

Verify antiemetics taken or administer as prescribed.

## ⌚ Chemotherapy - Time out

### Calcium Folate (Leucovorin)

- administer via IV infusion over 2 hours
- flush with ~ 50 mL of sodium chloride 0.9%.

### Fluorouracil

- If using cryotherapy commence ~ 5 minutes prior to administering fluorouracil and continue for ~ 30 minutes post.

#### Administer fluorouracil (irritant):

- over 3 to 5 minutes
  - via a minibag **OR**
  - by IV bolus via a side port of a freely flowing IV infusion
- flush with ~ 100 mL of sodium chloride 0.9%.

### Fluorouracil continuous infusion (irritant)

Connect pump containing fluorouracil and administer over the correct time for the amount of drug in the pump:

- **▲ safety alert** issued for administration of infusional fluorouracil
- verify the correct rate of infusion via the ambulatory infusion pump
- read more information about the different [ambulatory infusion pumps](#).

Continue [safe handling](#) precautions until 7 days after completion of drug(s)

---

## Day 2

**Approximate treatment time: 2.5 hours**

[Safe handling and waste management](#)

[Safe administration](#)

[General patient assessment](#) prior to each treatment.

Any toxicity grade 2 or greater may require dose reduction, delay or omission of treatment and review by medical officer before recommencing treatment.

Stop ambulatory infusion pump.

Prime IV line(s).

### Pre treatment medication

Verify antiemetics taken or administer as prescribed.

## ⌚ Chemotherapy - Time out

### Calcium Folate (Leucovorin)

- administer via IV infusion over 2 hours
- flush with ~ 50 mL of sodium chloride 0.9%.

### Fluorouracil

- If using cryotherapy commence ~ 5 minutes prior to administering fluorouracil and continue for ~ 30 minutes post.

#### Administer fluorouracil (irritant):

- over 3 to 5 minutes
  - via a minibag **OR**
  - by IV bolus via a side port of a freely flowing IV infusion
- flush with ~ 100 mL of sodium chloride 0.9%.

### Fluorouracil (irritant)

Resume the continuous fluorouracil infusion:

- **▲ safety alert** issued for administration of infusional fluorouracil
- verify the correct rate of infusion via the ambulatory infusion pump.

Continue **safe handling** precautions until 7 days after completion of drug(s)

---

## Day 3

Approximate treatment time: 30 minutes

### Safe handling and waste management

### Disconnection of ambulatory infusion pump/infusor

Verify the ambulatory infusion pump/infusor is complete.

Disconnect the ambulatory infusion pump/infusor as per recommended procedure for type of pump/infusor.

Read more about [ambulatory infusion pumps/infusors](#).

Deaccess [TIVAD](#) or [CVAD](#).

Continue **safe handling** precautions until 7 days after completion of drug(s)

---

## Discharge information

### Antiemetics

- Antiemetics as prescribed.

### Antidiarrhoeals

- Antidiarrhoeals as prescribed.

### Patient information

- Ensure patient receives patient information sheet.

### Infusion pumps

- CADD-Legacy® 1 ambulatory infusion pump [patient information sheet](#).
- CADD-Legacy® Plus ambulatory infusion pump [patient information sheet](#).
- CADD® Solis VIP ambulatory infusion pump [patient information sheet](#).
- Elastomeric infusion system [patient information sheet](#).

## Side effects



The side effects listed below are not a complete list of all possible side effects for this treatment. Side effects are categorised into the approximate onset of presentation and should only be used as a guide.

Immediate (onset hours to days)	
<b>Nausea and vomiting</b>	Read more about <a href="#">prevention of treatment induced nausea and vomiting</a>
<b>Cardiotoxicity</b>	Coronary artery spasm is a temporary, sudden narrowing of one of the coronary arteries that may present at any time during treatment with fluoropyrimidines. It most commonly manifests as angina.
<b>Taste and smell alteration</b>	Read more about <a href="#">taste and smell changes</a>
Early (onset days to weeks)	
<b>Neutropenia</b>	Abnormally low levels of neutrophils in the blood. This increases the risk of infection. Any fever or suspicion of infection should be investigated immediately and managed aggressively. Read more about <a href="#">immediate management of neutropenic fever</a>
<b>Thrombocytopenia</b>	A reduction in the normal levels of functional platelets, increasing the risk of abnormal bleeding.  Read more about <a href="#">thrombocytopenia</a>
<b>Diarrhoea</b>	Read more about <a href="#">treatment induced diarrhoea</a>
<b>Ocular changes</b>	Symptoms may include eye pain, blurred vision, blepharitis, uveitis, optic neuritis, tear duct stenosis, conjunctivitis, hyperlacrimation, watery or dry eyes and photophobia.
<b>Oral mucositis</b>	Erythematous and ulcerative lesions of the gastrointestinal tract (GIT). It commonly develops following chemotherapy, radiation therapy to the head, neck or oesophagus, and high dose chemotherapy followed by a blood and marrow transplant (BMT). Read more about <a href="#">oral mucositis</a>
<b>Photosensitivity</b>	Increased sensitivity to ultraviolet (UV) light resulting in an exaggerated sunburn-like reaction accompanied by stinging sensations and urticaria.
<b>Actinic keratoses flare</b>	Pre-existing actinic keratoses (AKs) can become more inflamed and scaly as a result of immunosuppression. Read more about <a href="#">actinic keratoses flare</a>
<b>Fatigue</b>	Read more about <a href="#">fatigue</a>
<b>Palmar-plantar erythrodysesthesia (PPE) - hand-foot syndrome (HFS)</b>	Bilateral erythema, tenderness, pain, swelling, tingling, numbness, pruritus, dry rash, or moist desquamation and ulceration of the palms and soles. It is also known as hand-foot syndrome (HFS). Symptoms appear to be dose dependent and palms are affected more than soles. Read more about <a href="#">hand-foot syndrome associated with chemotherapy</a>
Late (onset weeks to months)	
<b>Anaemia</b>	Abnormally low levels of red blood cells (RBCs) or haemoglobin in the blood. Read more about <a href="#">anaemia</a>
<b>Alopecia - partial</b>	Hair thinning and/or patchy hair loss. Patients can also experience mild to moderate discomfort of the hair follicles, and rarely pain as the hair is falling out. Read more about <a href="#">alopecia</a> and <a href="#">scalp cooling</a>
<b>Hyperpigmentation</b>	Darkening of an area of skin caused by the overproduction of melanin.
<b>Nail changes</b>	Hyperpigmentation, paronychia, onycholysis, splinter haemorrhage, pyogenic granuloma formation, subungal haematoma and subungal hyperkeratosis are some of the nail changes associated with anti-cancer drugs. Read more about <a href="#">nail toxicities</a>

## Evidence

The evidence for using this regimen comes from a randomised 2 x 2 factorial designed trial of monthly fluorouracil plus leucovorin (FULV) also known as the Mayo regimen versus a bi-monthly fluorouracil plus leucovorin (LVFU2) also known as the De Gramont regimen for 24 or 36 weeks.<sup>1</sup>

Between July 1996 and November 1999, a total of 905 patients were enrolled and randomised to receive either:

1. Leucovorin 200mg/m<sup>2</sup> (IV infusion) plus 5-FU 400mg/m<sup>2</sup> (IV bolus) followed by 5-FU 600mg/m<sup>2</sup> (continuous infusion) for 22 hours for 2 consecutive days every 14 days or
2. Leucovorin 200mg/m<sup>2</sup> (IV infusion) plus 5-FU 400mg/m<sup>2</sup> (IV bolus) for 5 consecutive days every 28 days

In each arm the patients were then assigned to receive either 24- or 36 weeks of treatment.


The end points of the study were to compare the therapeutic ratio, efficacy, and toxicity of both regimens.<sup>1</sup>

### Efficacy


At a median follow up of 41 months, the disease-free survival (DFS) was similar between LVFU2 and FULV groups (HR=1.04;p=0.74) and between 24 and 36 weeks of therapy (HR=0.94;p=0.63).

Analysis of overall survival showed a slight excess, but insignificant, in the number of deaths in LVFU2 compared with FULV (HR=1.26;p=0.18).

#### Disease-free survival according to treatment<sup>1</sup>



#### Overall survival according to treatment<sup>1</sup>



## Toxicity

The most commonly observed grade 3 to 4 toxicities were neutropenia, diarrhoea and mucositis. Toxicities were significantly lower in the LVFU2 group (all toxicities,  $p < 0.001$ ).

### Maximum grade 3/4 toxicities per patient<sup>1</sup>

Table 4. Maximum Grade 3/4 Toxicities per Patient (National Cancer Institute common toxicity criteria)

	LVFU2, % of Patients (n = 452)	FULV, % of Patients (n = 453)	P*
Neutropenia	7	16	< .001
Diarrhea	4	9	< .001
Mucositis	2	7	< .001
Nausea/vomiting	1	3	.093
Hand-foot syndrome	0	0	—
All toxicities	11	26	< .001

Abbreviations: LVFU2, fluorouracil and leucovorin semimonthly; FULV, fluorouracil and leucovorin monthly.

\*Fisher's exact test.

© Journal of Clinical Oncology 2003

## References

- 1 Andre, T., P. Colin, C. Louvet, et al. 2003. "Semimonthly versus monthly regimen of fluorouracil and leucovorin administered for 24 or 36 weeks as adjuvant therapy in stage II and III colon cancer: results of a randomized trial." *J.Clin Oncol* 21(15):2896-2903.

## History

### Version 5

Date	Summary of changes
08/10/2019	Clinical information updated with PBS expanded indications for GCSF.
22/10/2020	Protocol reviewed electronically by Medical Oncology Reference Committee. No changes. Next review in 2 years.
20/10/2022	Protocol reviewed electronically by Medical Oncology Reference Committee. No changes. Next review in 4 years.

**Version 4**

Date	Summary of changes
01/05/2007	Patient sheet updated.
29/08/2007	Fluorouracil infusion reworded to total dose over 44 hours.
03/12/2007	BWI information and safety alert added.
20/08/2009	Review, new dose modifications and transferred to eviQ.
01/07/2010	Haematological dose modifications updated ( 20% changed to 25% dose reduction; cut-off for platelets for dose reduction changed from $10 \times 10^9/L$ to $50 \times 10^9/L$ ).
25/01/2011	New format to allow for export of protocol information. Protocol version number changed to V.2. Antiemetics and premedications added to the treatment schedule. Additional Clinical Information, Key Prescribing table and Key Administration table combined into new section titled Clinical Considerations. Drug specific information placed behind the drug name link.
15/3/2012	PHC view updated.
30/09/2013	Protocol reviewed at Medical Oncology Reference Committee. Group consensus to supersede protocol as modified degramont ID 76 is widely accepted as the preferred regimen. Next review in 2 years.
24/08/2014	PHC view removed.
18/02/2016	Discussion with Medical Oncology Reference Committee Chairs and protocol to be reviewed every 5 years. Next review due in 3 years.
16/10/2016	Patient information sheet updated to include more fluorouracil toxicity symptom warnings.
09/11/2016	The following changes made post Medical Oncology Reference Committee meeting held on 21 October 2016. Link to AGTIG and ANZCTR added. Sentence in dose modifications regarding omitting leucovorin if fluorouracil is delayed or omitted changed to specify fluorouracil bolus.
31/05/2017	Transferred to new eviQ website. Version number change to V.3.
16/02/2018	Protocol reviewed electronically by Medical Oncology Reference Committee. Fluoropyrimidine warning added. Review in 5 years.
10/05/2018	Haematological dose modifications updated as per consensus of the expert clinician group. Version number changed to V.4.

The information contained in this protocol is based on the highest level of available evidence and consensus of the eviQ reference committee regarding their views of currently accepted approaches to treatment. Any clinician (medical oncologist, haematologist, radiation oncologist, medical physicist, radiation therapist, pharmacist or nurse) seeking to apply or consult this protocol is expected to use independent clinical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use is subject to eviQ's disclaimer available at [www.eviq.org.au](http://www.eviq.org.au)

**First approved:** 14 November 2005  
**Last reviewed:** 20 October 2022  
**Review due:** 31 December 2026  
**Superseded:** 30 September 2013

***The currency of this information is guaranteed only up until the date of printing, for any updates please check:***

<https://www.eviq.org.au/p/75>

16 Jul 2023

# Patient information - Bowel cancer adjuvant - de Gramont (fluorouracil and leucovorin)

Patient's name:

---

## Your treatment

The treatment schedule below explains how the drugs for this treatment are given.


### de Gramont (fluorouracil and leucovorin)

This treatment cycle is repeated every 14 days. You will have 12 cycles.

Day	Treatment	How it is given	How long it takes
1	<b>Calcium folinate (Leucovorin)</b> <i>(loo-koe-VOR-in)</i>	By a drip into a vein	About 2.5 hours
	<b>Fluorouracil</b> <i>(flure-oh-YOOR-a-sill)</i>	By a drip into a vein and then slowly through a pump	For 22 hours by pump at home
2	<b>Calcium folinate (Leucovorin)</b>	By a drip into a vein	About 2.5 hours
	<b>Fluorouracil</b>	By a drip into a vein and then slowly through a pump	For 22 hours by pump at home
3	<b>Disconnect pump</b>		About 30 minutes

## When to get help

Anticancer drugs (drugs used to treat cancer) can sometimes cause serious problems. It is important to get medical help immediately if you become unwell.

 <p><b>IMMEDIATELY go to your nearest hospital Emergency Department, or contact your doctor or nurse if you have any of the following at any time:</b></p>	<p><b>Emergency contact details</b></p> <p>Ask your doctor or nurse from your treating team who to contact if you have a problem</p>
<ul style="list-style-type: none"> <li>• a temperature of 38°C or higher</li> <li>• chills, sweats, shivers or shakes</li> <li>• shortness of breath</li> <li>• uncontrolled vomiting or diarrhoea</li> <li>• pain, tingling or discomfort in your chest or arms</li> <li>• leaking from your pump</li> <li>• you become unwell.</li> </ul>	<p>Daytime:.....</p> <p>Night/weekend:.....</p> <p>Other instructions:.....</p> <p>.....</p> <p>.....</p> <p>.....</p>

**During your treatment immediately** tell the doctor or nurse looking after you if you get any of the following problems:

- leaking from the area where the drugs are being given
- pain, stinging, swelling or redness in the area where the drugs are being given or at any injection sites
- a skin rash, itching, feeling short of breath, wheezing, fever, shivers, or feeling dizzy or unwell in any way (allergic reaction).

### Changes to your dose or treatment delays

Sometimes a treatment may be started at a lower dose or the dose needs to be changed during treatment. There may also be times when your treatment is delayed. This can happen if your doctor thinks you are likely to have severe side effects, if you get severe side effects, if your blood counts are affected and causing delays in treatment, or if you are finding it hard to cope with the treatment. This is called a dose reduction, dose change or treatment delay. Your doctor will explain if you need any changes or delays to your treatment and the reason why.

### Blood tests and monitoring

Anti-cancer drugs can reduce the number of blood cells in your body. You will need to have regular blood tests to check that your blood cell count has returned to normal. If your blood count is low your treatment may be delayed until it has returned to normal. Your doctor or nurse will tell you when to have these blood tests. Tell your doctor if you are on an anticoagulant (medication used to treat or prevent blood clots) e.g. warfarin. You may need to have additional blood tests.

### Pumps and central venous access devices (CVADs)

This treatment involves having chemotherapy through a pump. To have this, you will also need a central venous access device (CVAD). Your doctor or nurse will explain this to you. For more information see the [eviQ patient information sheets on pumps and CVADs](#). At home you will need to look at your pump 3 to 4 times a day to check it is working. Your nurse will teach you how to do this.

### Other medications given during this treatment

- **Anti-sickness (anti-nausea) medication:** you may be given some anti-sickness medication. Make sure you take this medication as your doctor or nurse tells you, even if you don't feel sick. This can help to prevent the sickness starting.
- **Antidiarrhoeals:** you may be given some medication to treat diarrhoea. Your doctor or nurse will tell you how and when to take your antidiarrhoeal medication.

### Superseded treatments

This treatment is superseded meaning that better treatments have taken its place. Uncommonly superseded treatments are still used. Your doctor will explain why this treatment has been selected for you.

## Side effects

Cancer treatments can cause damage to normal cells in your body, which can cause side effects. Everyone gets different side effects, and some people will have more problems than others.

The table below shows some of the side effects you may get with this treatment. You are unlikely to get all of those listed and you may also get some side effects that have not been listed.

Tell your doctor or nurse about any side effects that worry you. Follow the instructions below and those given to you by your doctor or nurse.

Immediate (onset hours to days)	
<b>Nausea and vomiting</b>	<ul style="list-style-type: none"> <li>You may feel sick (nausea) or be sick (vomit).</li> <li>Take your anti-sickness medication as directed even if you don't feel sick.</li> <li>Drink plenty of fluids (unless you are fluid restricted).</li> <li>Eat small meals more frequently.</li> <li>Try food that does not require much preparation.</li> <li>Try bland foods like dry biscuits or toast.</li> <li>Gentle exercise may help with nausea.</li> <li>Ask your doctor or nurse for eviQ patient information - <a href="#">Nausea and vomiting during cancer treatment</a>.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have uncontrolled vomiting or feel dizzy or light-headed.</b></li> </ul>
<b>Heart problems</b>	<ul style="list-style-type: none"> <li>You may get: <ul style="list-style-type: none"> <li>chest pain or tightness</li> <li>shortness of breath</li> <li>an abnormal heartbeat</li> </ul> </li> <li>Tell your doctor if you have a history of heart problems or high blood pressure.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the symptoms listed above.</b></li> </ul>
<b>Taste and smell changes</b>	<ul style="list-style-type: none"> <li>You may find that food loses its taste or tastes different.</li> <li>These changes are likely to go away with time.</li> <li>Do your mouth care regularly.</li> <li>Chew on sugar-free gum or eat sugar-free mints.</li> <li>Add flavour to your food with sauces and herbs.</li> <li>Ask your doctor or nurse for eviQ patient information - <a href="#">Taste and smell changes during cancer treatment</a>.</li> </ul>

Early (onset days to weeks)	
<b>Infection risk (neutropenia)</b>	<ul style="list-style-type: none"> <li>This treatment lowers the amount of white blood cells in your body. The type of white blood cells that help to fight infection are called neutrophils. Having low level of neutrophils is called neutropenia. If you have neutropenia, you are at greater risk of getting an infection. It also means that your body can't fight infections as well as usual. This is a serious side effect, and can be life threatening.</li> <li>Wash your hands often.</li> <li>Keep a thermometer at home and take your temperature regularly, and if you feel unwell.</li> <li>Do your mouth care regularly.</li> <li>Inspect your central line site (if you have one) daily for any redness, pus or swelling.</li> <li>Limit contact with people who are sick.</li> <li>Learn how to recognise the signs of infection.</li> <li>Ask your doctor or nurse for eviQ patient information - <a href="#">Infection during cancer treatment</a>.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you get any of the following signs or symptoms:</b> <ul style="list-style-type: none"> <li><b>a temperature of 38°C or higher</b></li> <li><b>chills, shivers, sweats or shakes</b></li> <li><b>a sore throat or cough</b></li> <li><b>uncontrolled diarrhoea</b></li> <li><b>shortness of breath</b></li> <li><b>a fast heartbeat</b></li> <li><b>become unwell even without a temperature.</b></li> </ul> </li> </ul>

<p><b>Low platelets (thrombocytopenia)</b></p>	<ul style="list-style-type: none"> <li>• This treatment lowers the amount of platelets in your blood. Platelets help your blood to clot. When they are low, you are at an increased risk of bleeding and bruising.</li> <li>• Try not to bruise or cut yourself.</li> <li>• Avoid contact sport or vigorous exercise.</li> <li>• Clear your nose by blowing gently.</li> <li>• Avoid constipation.</li> <li>• Brush your teeth with a soft toothbrush.</li> <li>• Don't take aspirin, ibuprofen or other similar anti-inflammatory medications unless your doctor tells you to.</li> <li>• Tell your doctor or nurse if you have any bruising or bleeding.</li> <li>• <b>Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if you have any uncontrolled bleeding.</b></li> </ul>
<p><b>Diarrhoea</b></p>	<ul style="list-style-type: none"> <li>• You may get bowel motions (stools, poo) that are more frequent or more liquid.</li> <li>• You may also get bloating, cramping or pain.</li> <li>• Take your antidiarrhoeal medication as directed by your doctor.</li> <li>• Drink plenty of fluids (unless you are fluid restricted).</li> <li>• Eat and drink small amounts more often.</li> <li>• Avoid spicy foods, dairy products, high fibre foods, and coffee.</li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Diarrhoea during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse immediately, or go to your nearest hospital Emergency Department if your diarrhoea is not controlled, you have 4 or more loose bowel motions per day, and if you feel dizzy or light-headed.</b></li> </ul>
<p><b>Eye problems</b></p>	<ul style="list-style-type: none"> <li>• You may get: <ul style="list-style-type: none"> <li>◦ eye pain</li> <li>◦ red, sore or swollen eyes</li> <li>◦ blurred vision</li> <li>◦ watery or gritty eyes</li> <li>◦ changes in your eyesight</li> <li>◦ sensitivity to sunlight.</li> </ul> </li> <li>• Protect your eyes from the weather (sun and wind) by wearing sunglasses, especially if you have lost your eyelashes.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above. Eye drops may help with your symptoms.</b></li> </ul>
<p><b>Mouth pain and soreness (mucositis)</b></p>	<ul style="list-style-type: none"> <li>• You may have: <ul style="list-style-type: none"> <li>◦ bleeding gums</li> <li>◦ mouth ulcers</li> <li>◦ a white coating on your tongue</li> <li>◦ pain in the mouth or throat</li> <li>◦ difficulty eating or swallowing.</li> </ul> </li> <li>• Avoid spicy, acidic or crunchy foods and very hot or cold food and drinks.</li> <li>• Try bland and soft foods.</li> <li>• Brush your teeth gently with a soft toothbrush after each meal and at bedtime. If you normally floss continue to do so.</li> <li>• Rinse your mouth after you eat and brush your teeth, using either: <ul style="list-style-type: none"> <li>◦ 1/4 teaspoon of salt in 1 cup of warm water, or</li> <li>◦ 1/4 teaspoon of bicarbonate of soda in 1 cup of warm water</li> </ul> </li> <li>• Ask your doctor or nurse for eviQ patient information - <a href="#">Mouth problems during cancer treatment</a>.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>
<p><b>Skin that is more sensitive to the sun (photosensitivity)</b></p>	<ul style="list-style-type: none"> <li>• After being out in the sun you may develop a rash like a bad sunburn.</li> <li>• Your skin may become red, swollen and blistered.</li> <li>• Avoid direct sunlight.</li> <li>• Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and a sunscreen of SPF 50 or higher.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>



<b>Skin changes</b>	<ul style="list-style-type: none"> <li>• Your skin may become dry, and you may notice changes to areas of your skin that have been exposed to the sun.</li> <li>• Keep your skin moisturised with a cream such as sorbolene or aqueous cream.</li> <li>• Avoid direct sunlight.</li> <li>• Protect your skin from the sun by wearing a wide-brimmed hat, sun-protective clothing, sunglasses and sunscreen of SPF 50 or higher.</li> <li>• <b>Tell your doctor or nurse if you notice any skin changes.</b></li> </ul>
<b>Tiredness and lack of energy (fatigue)</b>	<ul style="list-style-type: none"> <li>• You may feel very tired, have no energy, sleep a lot, and not be able to do normal activities or things you enjoy.</li> <li>• Do not drive or operate machinery if you are feeling tired.</li> <li>• Nap for short periods (only 1 hour at a time)</li> <li>• Prioritise your tasks to ensure the best use of your energy.</li> <li>• Eat a well balanced diet and drink plenty of fluids (unless you are fluid restricted).</li> <li>• Try some gentle exercise daily.</li> <li>• Allow your friends and family to help.</li> <li>• <b>Tell your doctor or nurse if you get any of the symptoms listed above.</b></li> </ul>
<b>Hand-foot syndrome (palmar-plantar erythrodysesthesia)</b>	<ul style="list-style-type: none"> <li>• The palms of your hands and soles of your feet may become: <ul style="list-style-type: none"> <li>◦ red and hot</li> <li>◦ swollen</li> <li>◦ painful and tender</li> <li>◦ blistered.</li> </ul> </li> <li>• The skin in the area may also peel.</li> <li>• Moisturise your hands and feet daily with sorbolene or aqueous cream.</li> <li>• Keep your hands and feet clean and dry.</li> <li>• Avoid hot water, instead use lukewarm water to bathe.</li> <li>• Avoid direct sunlight.</li> <li>• Avoid unnecessary walking, jogging or exercise.</li> <li>• Wear cotton socks and avoid tight-fitting shoes.</li> <li>• <b>Tell your doctor or nurse as soon as possible if you notice any skin changes on your hands or feet.</b></li> </ul>

Late (onset weeks to months)	
<b>Low red blood cells (anaemia)</b>	<ul style="list-style-type: none"> <li>You may feel dizzy, light-headed, tired and appear more pale than usual.</li> <li>Tell your doctor or nurse if you have any of these signs or symptoms. You might need a blood transfusion.</li> <li><b>Tell your doctor or nurse immediately, or go to the nearest hospital Emergency Department if you have any chest pain, trouble breathing, or feel like your heart is racing.</b></li> </ul>
<b>Hair thinning</b>	<ul style="list-style-type: none"> <li>Your hair may become dry and may break easily.</li> <li>You may lose some of your hair.</li> <li>Use a gentle shampoo and a soft hairbrush.</li> <li>Take care with hair products like hairspray, hair dye, bleaches and perms.</li> <li>Protect your scalp from the cold with a hat or scarf.</li> <li>Protect your scalp from the sun with a hat and sunscreen of SPF 50 or higher.</li> <li>Ask your doctor or nurse about the <a href="http://www.lgfb.org.au">Look Good Feel Better</a> program (www.lgfb.org.au)</li> </ul>
<b>Skin colour changes</b>	<ul style="list-style-type: none"> <li>You may have darkening of your skin, especially in areas that are exposed to the sun.</li> <li>You may also notice darkening of your tongue, gums and over your finger joints.</li> <li>These skin changes may fade over time.</li> <li>Protect your skin from the sun by wearing sun-protective clothing, a wide-brimmed hat, sunglasses and a sunscreen of SPF 50 or higher.</li> </ul>
<b>Nail changes</b>	<ul style="list-style-type: none"> <li>Your nails may: <ul style="list-style-type: none"> <li>grow more slowly</li> <li>become darker</li> <li>develop ridges or white lines</li> <li>become brittle and flaky</li> </ul> </li> <li>In some cases, you may lose your nails completely.</li> <li>Keep your nails clean and short.</li> <li>Avoid things like biting your fingernails, getting a manicure, pedicure or false nails.</li> <li>Wear gloves when you wash the dishes, work in the garden, or clean the house.</li> </ul>

## General advice for people having cancer treatment

### Chemotherapy safety

- Learn how to keep you and your family safe while you are having anticancer drugs.
- See our patient information sheet - [Chemotherapy safety at home](#).

### Blood clot risk

- Cancer and anticancer drugs can increase the risk of a blood clot (thrombosis).
- Tell your doctor if you have a family history of blood clots.
- A blood clot can cause pain, redness, swelling in your arms or legs, shortness of breath or chest pain.
- If you have any of these symptoms go to your nearest hospital Emergency Department.

### Medications and vaccinations

- Before you start treatment, tell your doctor about any medications you are taking, including vitamins or herbal supplements.
- Don't stop or start any medications during treatment without talking to your doctor and pharmacist first.
- Paracetamol is safe to take if you have a headache or other mild aches and pains. It is recommended that you avoid taking aspirin, ibuprofen and other anti-inflammatory type medications for pain while you are having treatment. However, if these medications have been prescribed by your doctor, do not stop taking them without speaking with your doctor.
- Vaccinations such as flu and tetanus vaccines are safe to receive while having treatment. Do not have any live vaccines during your treatment or for 6 months after it finishes. If you are unsure, check with your doctor before you have any vaccinations.
- People you live with should be fully vaccinated, including having live vaccines according to the current vaccination schedule. Extra care needs to be taken with hand washing and careful disposal of soiled nappies for infants who have recently received the rotavirus vaccine.

## Other medical and dental treatment

- If you go to hospital or any other medical appointment (including dental appointments), always tell the person treating you that you are receiving anticancer drugs.
- Before you have any dental treatment, talk to your doctor.

## Diet

- While you are receiving this treatment it is important that you try to maintain a healthy diet.
- Speak to your doctor or nurse about whether drinking alcohol is safe with your treatment.
- If you have any concerns about recent weight loss or weight gain or questions about your diet, ask to speak to a dietitian.

## Fertility

- Some cancer treatments can reduce your fertility. This can make it difficult or impossible to get pregnant or father a child.
- Talk to your doctor or nurse before you start any treatment. Depending on your situation there may be fertility sparing options available to you and/or your partner, discuss these with your doctor or nurse.

## Pregnancy and breastfeeding

- Some cancer treatments can be dangerous to unborn babies. Talk to your doctor or nurse if you think there is any chance that you could be pregnant.
- Do not try to get pregnant or father a child during this treatment. Contraception should be used during treatment and after stopping treatment. Ask your doctor or nurse about what type of contraception you should use.
- If you are planning pregnancy/fatherhood after completing this treatment, talk to your doctor. Some doctors advise waiting between 6 months and 2 years after treatment.
- Do not breastfeed if you are on this treatment, as anti-cancer medications can also pass into breast milk.

## Sex life and sexuality

- The desire to have sex may decrease as a result of this treatment or its side effects.
- Your emotions and the way you feel about yourself may also be affected by this treatment.
- It may help to discuss your concerns with your partner and doctor or nurse.

## Quitting smoking

- It is never too late to quit smoking. Quitting smoking is one of the best things you can do to help your treatment work better.
- There are many effective tools to improve your chances of quitting.
- Talk to your treating team for more information and referral to a smoking cessation support service.

## Staying active

- Research shows that exercise, no matter how small, has many benefits for people during and after cancer treatment.
- Talk to your doctor before starting an exercise program. Your doctor can advise whether you need a modified exercise program.

**For more information about cancer treatment, side effects and side effect management see our [Patient and carers](#) section.**

## Where to get more information

### Telephone support

- Call Cancer Council on 13 11 20 for cancer information and support

### Bowel cancer information

- Australian Council of Stoma Associations – [australianstoma.com.au](http://australianstoma.com.au)
- Australian Government Bladder and Bowel – [bladderbowel.gov.au](http://bladderbowel.gov.au)
- Australian Government Department of Health & Ageing Stoma appliance scheme – [health.gov.au/internet/main/publishing.nsf/Content/Stoma+Appliance+Scheme-1](http://health.gov.au/internet/main/publishing.nsf/Content/Stoma+Appliance+Scheme-1)
- Bowel Cancer Australia – [bowelcanceraustralia.org](http://bowelcanceraustralia.org)
- National Public Toilet map – [toiletmap.gov.au](http://toiletmap.gov.au)
- Recovering after Pelvic Radiation Therapy: A guide for women – <https://www.targetingcancer.com.au/useful-resources/recovering-after-pelvic-radiation-therapy-a-guide-for-women/>

### General cancer information and support

- Australian Rare Cancer (ARC) Portal – [arcportal.org.au/](http://arcportal.org.au/)
- Beyondblue – [beyondblue.org.au](http://beyondblue.org.au)
- Cancer Australia – [canceraustralia.gov.au](http://canceraustralia.gov.au)
- Cancer Council Australia – [cancer.org.au](http://cancer.org.au)
- Cancer Voices Australia – [cancervoicesaustralia.org](http://cancervoicesaustralia.org)
- CanTeen – [canteen.org.au](http://canteen.org.au)
- Carers Australia – [carersaustralia.com.au](http://carersaustralia.com.au)
- CHILL Cancer related hair loss - [scalpcooling.org](http://scalpcooling.org)
- eviQ Cancer Treatments Online – [eviQ.org.au](http://eviQ.org.au)
- LGBTQI+ People and Cancer - [cancercouncil.com.au/cancer-information/lgbtqi](http://cancercouncil.com.au/cancer-information/lgbtqi)
- Look Good Feel Better – [lgfb.org.au](http://lgfb.org.au)
- Patient Information – [patients.cancer.nsw.gov.au](http://patients.cancer.nsw.gov.au)
- Radiation Oncology Targeting Cancer – [targetingcancer.com.au](http://targetingcancer.com.au)
- Redkite – [redkite.org.au](http://redkite.org.au)
- Return Unwanted Medicines – [returnmed.com.au](http://returnmed.com.au)
- Staying active during cancer treatment – [patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active](http://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/staying-active)

**Quit smoking information and support**

Quitting smoking is helpful even after you have been diagnosed with cancer. The following resources provide useful information and support to help you quit smoking. Talk to your treating team about any other questions you may have.

- Call Quitline on 13 QUIT (13 78 48)
- iCanQuit – [iCanQuit.com.au](http://iCanQuit.com.au)
- Patient Information - [patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking](http://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/quitting-smoking)
- Quitnow – [quitnow.gov.au](http://quitnow.gov.au)

**Additional notes:**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

This document is a guide only and cannot cover every possible situation. The health professionals caring for you should always consider your individual situation when making decisions about your care. Contact your cancer clinic staff or doctor if you have any questions or concerns about your treatment, or you are having problems coping with side effects. While eviQ endeavours to link to reliable sources that provide accurate information, eviQ and the Cancer Institute NSW do not endorse or accept responsibility for the accuracy, currency, reliability or correctness of the content of linked external information sources. Use of this document is subject to eviQ's disclaimer available at [www.eviq.org.au](http://www.eviq.org.au)

---

**First approved:** 14 November 2005

**Last reviewed:** 20 October 2022

**Review due:** 31 December 2026

**Superseded:** 30 September 2013

***The currency of this information is guaranteed only up until the date of printing, for any updates please check:***

*<https://www.eviq.org.au/pi/75>*

*16 Jul 2023*